

MicroRNA profiling of paediatric AML with *FLT-ITD* or *MLL*-rearrangements: Expression signatures and *in vitro* modulation of miR-221-3p and miR-222-3p with BRD4/HATs inhibitors

PIER PAOLO LEONCINI^{1*}, PATRIZIA VITULLO^{1*}, SOFIA REDDEL¹, VALERIA TOCCO¹, VALERIA PAGANELLI¹,
FRANCESCA STOCCHI¹, ELENA MARIGGIÒ¹, MICHELE MASSA², GIOVANNI NIGITA³,
DARIO VENEZIANO³, PAOLO FADDA⁴, MARIO SCARPA⁵, MARTINA PIGAZZI^{6,7},
ALICE BERTAINA¹, ROSSELLA ROTA¹, DARIA PAGLIARA^{1*} and PIETRO MERLI^{1*}

¹Department of Paediatric Haematology and Oncology, Cell and Gene Therapy, Bambino Gesù Children's Hospital, IRCCS, I-00146 Rome; ²Department of Maternal Infantile and Urological Sciences, Sapienza University of Rome, I-00161 Rome, Italy; ³Department of Cancer Biology and Genetics, Comprehensive Cancer Center; ⁴Genomics Shared Resource, Comprehensive Cancer Center, The Ohio State University, Columbus, OH 43210-1239, USA; ⁵Department of Biomedical Research, Urology Research Laboratory, University of Bern, CH-3008 Bern, Switzerland; ⁶Department of Women's and Children's Health (SDB), Hematology-Oncology Laboratory, University of Padova, I-35128 Padova; ⁷Pediatric Onco-Hematology, Stem Cell Transplant and Gene Therapy Laboratory, Istituto di Ricerca Pediatrica - Città della Speranza, I-35127 Padova, Italy

Received November 30, 2021; Accepted August 10, 2022

DOI: 10.3892/or.2022.8436

Abstract. Novel therapeutic strategies are needed for paediatric patients affected by Acute Myeloid Leukaemia (AML), particularly for those at high-risk for relapse. MicroRNAs (miRs) have been extensively studied as biomarkers in cancer and haematological disorders, and their expression has been correlated to the presence of recurrent molecular abnormalities, expression of oncogenes, as well as to prognosis/clinical outcome. In the present study, expression signatures of different miRs related both to presence of myeloid/lymphoid or mixed-lineage leukaemia 1 and Fms like tyrosine kinase 3 internal tandem duplications rearrangements and to the clinical outcome of paediatric patients with AML were identified. Notably, miR-221-3p and miR-222-3p resulted as a possible relapse-risk related miR. Thus, miR-221-3p and miR-222-3p expression modulation was investigated by using a Bromodomain-containing protein 4

(BRD4) inhibitor (JQ1) and a natural compound that acts as histone acetyl transferase inhibitor (curcumin), alone or in association, in order to decrease acetylation of histone tails and potentiate the effect of BRD4 inhibition. JQ1 modulates miR-221-3p and miR-222-3p expression in AML with a synergic effect when associated with curcumin. Moreover, changes were observed in the expression of *CDKN1B*, a known target of miR-221-3p and miR-222-3p, increase in apoptosis and down-regulation of miR-221-3p and miR-222-3p expression in CD34⁺ AML primary cells. Altogether, these findings suggested that several miRs expression signatures at diagnosis may be used for risk stratification and as relapse prediction biomarkers in paediatric AML outlining that epigenetic drugs, could represent a novel therapeutic strategy for high-risk paediatric patients with AML. For these epigenetic drugs, additional research for enhancing activity, bioavailability and safety is needed.

Introduction

Acute Myeloid Leukaemia (AML) constitutes 20% of all paediatric leukaemia and is responsible for substantial mortality. Despite progress made over the past years in diagnosis, risk stratification and treatment of AML, survival remains suboptimal with a success rate of 60-70% (1-3), with relapse being the leading cause of death. Risk stratification in patients with AML is of paramount importance in order to deliver tailored therapy, enabling treatment intensification in high-risk patients (for example, allogeneic stem cell transplantation in high-risk in first complete remission). Moreover, novel therapies are needed in order to further improve prognosis in these patients. Among

Correspondence to: Dr Sofia Reddel, Department of Paediatric Haematology and Oncology, Cell and Gene Therapy, Bambino Gesù Children's Hospital, IRCCS, 15 Viale di San Paolo, I-00146 Rome, Italy
E-mail: sofia.reddel@opbg.net; sofia.reddel@gmail.com

*Contributed equally

Key words: acute myelogenous leukaemia, microRNAs, epigenetic drugs, biomarkers, children

prognostic factors, cytogenetic and molecular abnormalities, together with Minimal Residual Disease and treatment response, play a pivotal role in defining AML prognosis and treatment (4,5). Rearrangements involving Histone-lysine N-methyltransferase 2A (*KMT2A*) gene, formerly known as myeloid/lymphoid or mixed-lineage leukaemia 1 (*MLL1*) gene, as well as Fms like tyrosine kinase 3 (*FLT3*) internal tandem duplications (*FLT3-ITD*) represent useful prognostic factors and, possibly, therapeutic targets. Indeed, they still define an intermediate to high risk AML (1-3,6). In particular, *FLT3-ITD* mutations are associated with higher risk of relapse and dismal prognosis (1,3,7), whereas *MLL*-rearranged AML is an heterogeneous group of diseases with more than 100 rearrangements being described and with different outcome largely dependent on the fusion partner (8).

FLT3-ITD and *MLL* rearrangements have been functionally linked to dysregulation of expression of microRNAs (miRNAs or miRs) (9). miRs are small non-coding RNA molecules (~18-22 nucleotides long), involved in several cellular processes (10,11). In cancer, they are implicated both in promoting carcinogenesis (oncomiRs) and in suppressing tumour transformation (12). Their role in AML have been extensively investigated over the past years reviling promising data on diagnosis, prognostic stratification and, possibly, treatment in AML patients (13-15). Despite extensive research performed to understand the role of miRs in AML, the majority of studies are focused on adult patients, while a precise characterization of miRNAs expression in paediatric AML is less documented. Moreover, data on the role of different miRs are conflicting because of the variability of genetic abnormalities found in AML (16). In the present study, it was aimed to identify AML specific miR signatures in a cohort of patients harbouring molecular lesions (*FLT3-ITD* and *MLL* rearrangement), studying the expression of distinct miR sets in relation to relapse risk.

Epigenetic networks, including histone modification mechanisms, are involved in the regulation of both miRs expression and function. An increasing interest in the field of cancer therapeutic drugs is focused on small molecular compounds targeting epigenetic regulation (17).

Bromodomain and extra-terminal domain family of proteins (BET) and histone acetyl transferase (HAT) inhibitors proteins are the best characterized ones. BET are effective in preventing Bromodomain Containing protein 4 (BRD4) associated transcription of several oncogenes, reducing proliferation and increasing apoptosis in AML (18-22). BRD4 is a member of BET family proteins, characterized by the presence of functional structures called bromodomains which bind specific acetylated residues on histone tails to modulate transcription of target gene (23-25), enhancing transcription of several oncogenes (26).

Among BRD4 inhibitors, JQ1 was used as its activity on modulation of miRs was previously described (27). JQ1 [(S)-tert-butyl-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)acetate] is a small molecule belonging to the thienotriazolodiazepine group and it prevents the binding of BRD4 to acetylated residues on histone H3 tails, particularly H3AcK14 (20,28-30).

Furthermore, among HAT inhibitors, curcumin, a natural compound extracted from the root of *Curcuma Longa* has been shown to inhibit acetylation of histone tails, blocking the activity

of the HAT p300 even causing its proteasomal degradation (31). This results in a global decrease of acetylation on histone tails and a consequent modulation of gene transcription (32,33).

BRD4 inhibitors have exhibited only moderate results in clinics and novel ways to increase their antitumour activity are needed (34). It was therefore hypothesized that the association with curcumin would increase JQ1 efficacy. The BET family are 'readers' of chromatin acetylation whereas HAT could be classified as a 'writer' of histone acetylation (34) thus offering a theoretical basis for JQ1 and curcumin synergic activity.

Moreover, it was previously showed that also JQ1, like curcumin, blocks p300-mediated acetylation (25,35). Thus, it was investigated *in vitro* whether a combination of BRD4 and HAT inhibitors have an effect in terms of modulation of miRs and antitumour effects on different AML cell-lines harbouring mutations resembling those present in our patients.

Materials and methods

Samples of patients. A total of 23 patients aged 1 to 18 years, who received a diagnosis of AML harbouring *FLT3-ITD* or *MLL* rearrangement (Table SI). Although not mutually exclusive, these rearrangements are not frequently found together. In the present study, none of the patients had both the rearrangements.

Bone marrow (BM) samples were collected from January 1st 2010 to December 31st 2016 at Bambino Gesù Children's Hospital in Rome and at Department of Paediatrics, University in Padua, at diagnosis and at disease recurrence from the 13 patients who underwent relapse (REL-D and REL-R groups, respectively) and at diagnosis from the 10 patients who did not display relapse (NREL group).

A total of 8 frozen age-matched BM samples from healthy children (HD) (unused aliquots from healthy BM donors) were retrieved from the tissue bank at Bambino Gesù Children's Hospital as a control population. Informed consent was obtained from either parents or legal guardians according to the Declaration of Helsinki (2008). The present study was approved by the Institutional Review Boards of Bambino Gesù Children's Hospital (Rome, Italy).

CD34⁺ cells isolation. Mononuclear cells were isolated by density gradient centrifugation at 400 g and 20°C for 30 min, diluted in 90% fetal bovine serum (FBS) plus 10% dimethyl sulfoxide (DMSO) and stored in liquid nitrogen. CD34⁺ cells from BM samples of three patients randomly selected in our cohort, were magnetically separated using MACS CD34⁺ microbead kit (Miltenyi Biotec GmbH). In particular, the molecular analysis of these patients revealed a *FLT3-ITD* with normal karyotype and two *MLL* rearrangements [t(9;11) and t(10;11)]. The identity of CD34 cells was validated by flow cytometry using FACSCantoII equipped with FACSDiva 6.1 CellQuest software (Becton, Dickinson and Company) using 20 µl of CD34 PerCP antibody (cat. no 340666; BD Biosciences) with an incubation of 30 min at 4°C.

RNA extraction and miR profiling. Total RNA was extracted using TRIzol[®] reagent (Invitrogen; Thermo Fisher Scientific, Inc.) and purified using RNA Cleanup and Concentration kit according to the manufacturer's protocol (Norgen Biotek

Corp.). RNA quantification was performed using Nanodrop 2000 at 260 nm wavelength (Thermo Fisher Scientific, Inc.) and RNA integrity and purity was assessed with RNA Bioanalyzer kit according to the manufacturer's protocol (Agilent Technologies, Inc.). miR expression profile was performed using the nCounter Human v2 miRNA Expression Assay and nCounter Nanostring platform according to manufacturer's protocol (NanoString Technologies).

Cell culture methods. AML cell lines THP-1 (*MLL-MLLT3*; *MLL-AF9*), MOLM-13 (*MLL-MLLT3*; *MLL-AF9*; *FLT3-ITD*) and MV-4-11 (*MLL-AFF1*; *MLL-AF4*; *FLT3-ITD*) were obtained from DSMZ and cultured at 37°C using RPMI-1640 medium (Euroclone SpA) supplemented with 10% FBS (Thermo Fisher Scientific, Inc.) and 1% Penicillin-Streptomycin (Thermo Fisher Scientific, Inc.). Mycoplasma testing was performed for the cell lines used. Cell lines were treated with 250 nM JQ1 and 10 μ M curcumin singularly alone or in association, for 48 h. JQ1 and Curcumin were obtained from Selleck Chemicals and resuspended in DMSO, following the manufacturer's protocol.

Western blot analysis. Whole-cell lysates were prepared with RIPA lysis buffer (Thermo Fisher Scientific, Inc.) supplemented with protease and phosphatase inhibitors (Thermo Fisher Scientific, Inc.). Cells were lysed by sonication, incubated for 30 min at 4°C and then obtained cells lysates were centrifuged at 13,000 x g for 20 min at 4°C. The protein concentration of the resulting supernatant was estimated by BCA assay. Then, 40 μ g of sample was separated on Criterion TGX Precast Gels 4-20% (BioRad Laboratories, Inc.) and transferred to Hybond ECL nitrocellulose membranes (Amersham; Cytiva). Membranes were blocked at room temperature for 1 h in 5% non-fat milk in Tris buffered saline and 0.05% Tween-20 (TBS-T). Membranes were incubated at 4°C overnight with rabbit polyclonal anti-human CDKN1B (1:500; cat. no. sc-528; Santa Cruz Biotechnology, Inc.) and 1 h at room temperature with rabbit monoclonal anti-human GAPDH (1:1,000; cat. no. D16H11; Cell Signaling Technology, Inc.) primary antibodies. After incubation they were washed three times in TBS-T, then incubated with HRP-labelled goat anti-rabbit (1:5,000; cat. no. sc-2004) and goat anti-mouse (1:5,000; cat. no. sc-2005; both from Santa Cruz Biotechnology, Inc.) IgG secondary antibodies, respectively at room temperature for 1 h. Subsequently, they were washed an additional three times with TBS-T and then developed with ECL reagent (Western Lightning Plus; PerkinElmer, Inc.).

miR quantitative (q)PCR. Expression levels of hsa-miR-221-5p, hsa-miR-222-5p and U6 were measured using TaqMan microRNA assays (cat. nos. 000524, 002276 and 001973; Thermo Fisher Scientific, Inc.). Reverse transcription (RT) primer, preformulated forward/reverse primer and MGB probes for each assay were provided by the manufacturer. The TaqMan MiR Reverse Transcription kit was used for cDNA synthesis from 10 ng total RNA template according to the manufacturer's protocol. QuantStudio 12K Flex Real Time PCR System (Thermo Fisher Scientific, Inc.) was used for qPCR reactions with the following conditions: Enzyme activation 95°C for 20 sec and 40 cycles of denaturation (95°C for

1 sec) and annealing/extension (60°C for 20 sec) steps. miRNA expression data were normalized to U6 using the $2^{-\Delta\Delta C_q}$ (36) method by the Relative Quantification module of Thermo Fisher Cloud Data Analysis Apps. At least two independent amplifications were performed for each probe on triplicate samples.

Apoptosis assay. Following treatment with 250 nM BRD4 and 10 μ M curcumin, cells were washed twice with ice cold PBS and stained for 15 min at room temperature in calcium-binding buffer with PE-conjugated Annexin V (AnnV) and 7-Aminoactinomycin D (7-AAD) using the AnnV apoptosis detection kit (BD Pharmingen; BD Biosciences) according to the manufacturer's recommendations. Samples were analysed within 1 h by a fluorescence-activated cell sorting using a FACSCantoII equipped with FACSDiva 6.1 CellQuest software (Becton, Dickinson and Company).

Bioinformatics and statistical analyses. MicroRNA profiling normalization was performed using the nSolver Analysis Software (NanoString Technologies) as recommended by NanoString. P-values were calculated using the LIMMA (v.3.46.0) package (37) from the Bioconductor R (v.4.0.5) project. The P-values were adjusted for multiple testing using the Benjamini and Hochberg method to control the False Discovery Rate. An independent normalization phase for each comparison was performed, considering only the samples present in such comparison (for example, NREL vs. HD). Then, the miRNA expression of the HD group was specifically normalized in the comparisons in which the HD group was taken into consideration. Validated targets of miRs were reported in Table I according to miRWalk 2.0 online software analysis (<http://mirwalk.umm.uni-heidelberg.de/>). One-way ANOVA and post hoc comparison using Tukey's HSD Post Hoc or Dunnett's test were performed using SPSS software v19 (IBM Corp.) and GraphPad Prism v6 (GraphPad Software, Inc.). The heatmap was generated by using GenePattern tool (38), with Euclidean and Spearman correlation distances in columns and rows, respectively. Venn diagrams were created using web tool (39).

CD34⁺ cell culture. CD34⁺ cells were cultured using MethoCult H4434 methylcellulose medium (Stem Cell Technologies) supplemented with 250 nM JQ1 and 10 μ M curcumin.

Results

NanoString miR profiling reveals different expression signatures based on patient clinical and molecular features. A miR profiling analysis was first performed to verify whether the two distinct molecular subsets of the cohort of our patients (*MLL* rearranged and *FLT3-ITD*) showed different miR expression fingerprints. Comparing both *MLL* rearranged and *FLT3-ITD* sets with healthy donors (HDs), 4 and 16 significantly deregulated miRs were identified, respectively (Table I). miR-196b-5p and miR-34a-5p resulted upregulated in both AML sets. The comparison between the two molecular AML sets showed 3 differentially regulated miRs with miR-10a-5p and miR-99a-5p significantly higher in *FLT3-ITD* and miR-9a-5p with enhanced expression in *MLL*-rearranged sets (Table I). Other miRs

Table I. Molecular based miR expression signature.

FLT3-ITD vs. HD	Accession number	FC	Adjusted P-value
hsa-miR-10a-5p	MIMAT0000253	24.38	2x10 ⁻⁵
hsa-miR-451a	MIMAT0001631	17.06	1x10 ⁻³
hsa-miR-196b-5p	MIMAT0001080	4.31	5x10 ⁻⁴
hsa-miR-34a-5p	MIMAT0000255	3.38	4x10 ⁻³
hsa-let-7b-5p	MIMAT0000063	2.49	1x10 ⁻²
hsa-miR-29c-3p	MIMAT0000681	2.12	1x10 ⁻²
hsa-miR-520f-3p	MIMAT0002830	-2.11	4x10 ⁻²
hsa-miR-200c-3p	MIMAT0000617	-2.16	4x10 ⁻²
hsa-miR-421	MIMAT0003339	-2.34	2x10 ⁻²
hsa-miR-6724-5p	MIMAT0025856	-2.39	1x10 ⁻²
hsa-miR-151a-3p	MIMAT0000757	-2.45	1x10 ⁻²
hsa-miR-4755-5p	MIMAT0019895	-2.45	3x10 ⁻²
hsa-miR-365a-3p + hsa-miR-365b-3p	MIMAT0000710	-2.52	2x10 ⁻²
hsa-miR-520d-5p + hsa-miR-527 + hsa-miR-518a-5p	MIMAT0002855	-2.73	4x10 ⁻³
hsa-miR-574-5p	MIMAT0004795	-2.75	1x10 ⁻²
hsa-miR-342-3p	MIMAT0000753	-3.99	4x10 ⁻³
FLT3-ITD vs. MLL	Accession number	FC	Adjusted P-value
hsa-miR-10a-5p	MIMAT0000253	34.87	1.44x10 ⁻⁷
hsa-miR-99a-5p	MIMAT0000097	2.87	2.24x10 ⁻²
hsa-miR-9-5p	MIMAT0000441	-7.89	1.38x10 ⁻³
MLL vs. HD	Accession number	FC	Adjusted P-value
hsa-miR-9-5p	MIMAT0000441	10.75	9.33x10 ⁻³
hsa-miR-34a-5p	MIMAT0000255	5.50	8.78x10 ⁻⁵
hsa-miR-196b-5p	MIMAT0001080	4.23	1.55x10 ⁻²
hsa-miR-192-5p	MIMAT0000222	-2.73	3.92x10 ⁻²

hsa, *homo sapiens*; miR, microRNA; FLT3-ITD; Fms like tyrosine kinase 3 (*FLT3*) internal tandem duplications; HD, healthy donors.

such as miR-451a, miR-520d-5p/527/518a-5p, miR-574-5p and miR-192-5p were uniquely dysregulated in *FLT3-ITD* or *MLL* sets with respect to HDs (Table I). A miR expression profiling analysis was then performed based onto clinical outcomes of patients to identify those miRs associated with relapse.

The hierarchical clustering results of miRNAs expression of REL-D vs. NREL groups is revealed in Fig. 1. A total of 18 miRs were broadly upregulated in the REL-D patients set with respect to the NREL set and 48 miRs displayed the opposite trend (Fig. 1A and Table II). To further identify and refine a signature that was associated with relapse, among these differentially expressed 66 miRs, only those shared in the REL-D vs. HD and NREL vs. HD comparison were subsequently considered (Fig. 1B and Table II). The resulted signature associated to relapse and not-relapse is listed in Table III. Validated targets of miRs were identified using miRWalk 2.0 online software (40) and are reported in Table III. CDKN1B, a key regulator of cell cycle which has been previously reported to be associated with

prognosis in AML, resulted as primary target of miR-221-3p and miR-222-3p (41). To evaluate if the signature of miRs associated with relapse at diagnosis was maintained over time (if it is present also in the REL-R group), expression of miRs at diagnosis and at disease recurrence was compared between those patients who relapsed (REL-R vs. REL-D) and no significant differences were detected (Table II). Since our goal was to identify miRs associated to relapse with a significant prognostic value, the expression level of both miRs resulted overexpressed in REL-D vs. HD group and REL-D vs. NREL was first evaluated in the AML cells lines by qPCR (data not shown). High variability was obtained in the different cell lines. This result prompted the authors to focus only on hsa-miR-221-3p and hsa-miR-222-3p, presenting the same trend of expression in all the cell lines, as well as *in vivo* in the patients.

Association of JQ1 and curcumin leads to miR-221-3p and miR-222-3p downregulation and CDKN1B upregulation in

Table II. Clinical outcome-based miR expression signature.

REL-D vs. NREL	Accession number	FC	Adjusted P-value
hsa-miR-128-3p	MIMAT0000424	8,10	4x10 ⁻⁶
hsa-miR-143-3p	MIMAT0000435	7,47	9x10 ⁻⁴
hsa-miR-142-3p	MIMAT0000434	5,63	9x10 ⁻⁷
hsa-miR-144-3p	MIMAT0000436	3,87	4x10 ⁻²
hsa-miR-451a	MIMAT0001631	3,66	3x10 ⁻²
hsa-miR-29c-3p	MIMAT0000681	3,24	4x10 ⁻⁶
hsa-miR-140-5p	MIMAT0000431	2,53	2x10 ⁻⁵
hsa-miR-196b-5p	MIMAT0001080	2,48	7x10 ⁻⁴
hsa-miR-32-5p	MIMAT0000090	2,44	6x10 ⁻³
hsa-miR-221-3p	MIMAT0000278	2,04	1x10 ⁻²
hsa-miR-92a-3p	MIMAT0000092	1,93	4x10 ⁻³
hsa-miR-222-3p	MIMAT0000279	1,92	5x10 ⁻³
hsa-miR-21-5p	MIMAT0000076	1,88	1x10 ⁻²
hsa-miR-324-5p	MIMAT0000761	1,84	5x10 ⁻³
hsa-miR-34a-5p	MIMAT0000255	1,77	2x10 ⁻²
hsa-miR-30b-5p	MIMAT0000420	1,67	6x10 ⁻³
hsa-miR-30e-5p	MIMAT0000692	1,67	5x10 ⁻³
hsa-miR-106b-5p	MIMAT0000680	1,55	2x10 ⁻²
hsa-miR-30e-3p	MIMAT0000693	-1,41	3x10 ⁻²
hsa-miR-378i	MIMAT0019074	-1,44	4x10 ⁻²
hsa-miR-664a-3p	MIMAT0005949	-1,46	2x10 ⁻²
hsa-miR-24-3p	MIMAT0000080	-1,49	4x10 ⁻²
hsa-miR-25-3p	MIMAT0000081	-1,54	3x10 ⁻²
hsa-miR-2682-5p	MIMAT0013517	-1,67	6x10 ⁻²
hsa-miR-200c-3p	MIMAT0000617	-1,67	4x10 ⁻²
hsa-miR-320e	MIMAT0015072	-1,67	5x10 ⁻²
hsa-miR-191-5p	MIMAT0000440	-1,68	3x10 ⁻²
hsa-miR-423-5p	MIMAT0004748	-1,69	3x10 ⁻³
hsa-miR-362-5p	MIMAT0000705	-1,71	2x10 ⁻²
hsa-miR-149-5p	MIMAT0000450	-1,72	2x10 ⁻³
hsa-miR-194-5p	MIMAT0000460	-1,75	1x10 ⁻²
hsa-miR-6724-5p	MIMAT0025856	-1,80	5x10 ⁻³
hsa-miR-301a-3p	MIMAT0000688	-1,84	3x10 ⁻²
hsa-miR-378g	MIMAT0018937	-1,88	5x10 ⁻³
hsa-miR-132-3p	MIMAT0000426	-1,89	9x10 ⁻³
hsa-miR-28-3p	MIMAT0004502	-1,97	6x10 ⁻⁴
hsa-miR-28-5p	MIMAT0000085	-2,01	9x10 ⁻⁴
hsa-miR-574-3p	MIMAT0003239	-2,09	4x10 ⁻³
hsa-miR-520f-3p	MIMAT0002830	-2,10	5x10 ⁻⁵
hsa-miR-3144-3p	MIMAT0015015	-2,11	4x10 ⁻⁴
hsa-miR-1290	MIMAT0005880	-2,22	5x10 ⁻⁵
hsa-miR-421	MIMAT0003339	-2,24	6x10 ⁻⁴
hsa-miR-1275	MIMAT0005929	-2,32	2x10 ⁻⁴
hsa-miR-335-5p	MIMAT0000765	-2,39	5x10 ⁻⁴
hsa-miR-365a/b-3p	MIMAT0000710	-2,42	2x10 ⁻⁴
hsa-miR-625-5p	MIMAT0003294	-2,67	9x10 ⁻⁷
hsa-miR-612	MIMAT0003280	-2,70	2x10 ⁻⁶
hsa-miR-302d-3p	MIMAT0000718	-2,79	6x10 ⁻⁶
hsa-miR-150-5p	MIMAT0000451	-2,84	1x10 ⁻²
hsa-miR-301b-3p	MIMAT0004958	-2,90	8x10 ⁻⁷
hsa-miR-155-5p	MIMAT0000646	-2,95	3x10 ⁻³

Table II. Continued.

REL-D vs. NREL	Accession number	FC	Adjusted P-value
hsa-miR-556-5p	MIMAT0003220	-3,04	8x10 ⁻⁷
hsa-miR-3065-5p	MIMAT0015066	-3,07	3x10 ⁻⁷
hsa-miR-376a-3p	MIMAT0000729	-3,15	6x10 ⁻⁴
hsa-miR-342-3p	MIMAT0000753	-3,16	5x10 ⁻⁴
hsa-miR-197-5p	MIMAT0022691	-3,30	1x10 ⁻⁷
hsa-miR-4286	MIMAT0016916	-3,30	3x10 ⁻⁵
hsa-miR-585-3p	MIMAT0003250	-3,37	8x10 ⁻⁷
hsa-miR-642a-5p	MIMAT0003312	-3,41	5x10 ⁻³
hsa-miR-513a-3p	MIMAT0004777	-4,07	2x10 ⁻⁷
hsa-miR-1285-5p	MIMAT0022719	-4,10	1x10 ⁻⁷
hsa-miR-4284	MIMAT0016915	-4,82	8x10 ⁻⁷
hsa-miR-575	MIMAT0003240	-5,55	2x10 ⁻⁶
hsa-miR-1973	MIMAT0009448	-6,57	4x10 ⁻⁸
hsa-miR-3916	MIMAT0018190	-6,94	9x10 ⁻⁹
hsa-miR-378e	MIMAT0018927	-9,46	5x10 ⁻³
REL-D vs. HD	Accession number	FC	Adjusted P-value
hsa-miR-451a	MIMAT0001631	10,46	1x10 ⁻²
hsa-miR-196b-5p	MIMAT0001080	5,70	3x10 ⁻⁵
hsa-miR-34a-5p	MIMAT0000255	5,29	3x10 ⁻⁴
hsa-miR-221-3p	MIMAT0000278	2,14	2x10 ⁻²
hsa-miR-222-3p	MIMAT0000279	2,09	2x10 ⁻²
hsa-miR-29c-3p	MIMAT0000681	1,85	5x10 ⁻²
hsa-miR-941	MIMAT0004984	-1,87	5x10 ⁻²
hsa-miR-194-5p	MIMAT0000460	-1,94	5x10 ⁻²
hsa-miR-200c-3p	MIMAT0000617	-1,96	5x10 ⁻²
hsa-miR-192-5p	MIMAT0000222	-2,01	5x10 ⁻²
hsa-miR-1290	MIMAT0005880	-2,02	5x10 ⁻²
hsa-miR-612	MIMAT0003280	-2,08	5x10 ⁻²
hsa-miR-575	MIMAT0003240	-2,11	4x10 ⁻²
hsa-miR-151a-3p	MIMAT0000757	-2,11	4x10 ⁻²
hsa-miR-520f-3p	MIMAT0002830	-2,11	2x10 ⁻²
hsa-miR-421	MIMAT0003339	-2,16	3x10 ⁻²
hsa-miR-6724-5p	MIMAT0025856	-2,26	1x10 ⁻²
hsa-miR-4755-5p	MIMAT0019895	-2,30	3x10 ⁻²
hsa-miR-574-5p	MIMAT0004795	-2,32	3x10 ⁻²
hsa-miR-363-3p	MIMAT0000707	-2,38	3x10 ⁻²
hsa-miR-1285-5p	MIMAT0022719	-2,39	3x10 ⁻²
hsa-miR-4286	MIMAT0016916	-2,61	2x10 ⁻²
hsa-miR-365a/b-3p	MIMAT0000710	-3,00	3x10 ⁻³
hsa-miR-520d-5p/527/518a-5p	MIMAT0002855	-3,16	7x10 ⁻⁴
hsa-miR-342-3p	MIMAT0000753	-3,30	2x10 ⁻²
hsa-miR-150-5p	MIMAT0000451	-3,96	4x10 ⁻²
NREL vs. HD	Accession number	FC	Adjusted P-value
hsa-miR-9-5p	MIMAT0000441	6,90	8x10 ⁻³
hsa-miR-1973	MIMAT0009448	3,10	5x10 ⁻³
hsa-miR-4284	MIMAT0016915	2,82	3x10 ⁻²
hsa-miR-575	MIMAT0003240	2,72	5x10 ⁻²
hsa-miR-155-5p	MIMAT0000646	2,55	3x10 ⁻²

Table II. Continued.

NREL vs. HD	Accession number	FC	Adjusted P-value
hsa-miR-196b-5p	MIMAT0001080	2,37	5x10 ⁻²
hsa-miR-191-5p	MIMAT0000440	2,27	2x10 ⁻²
hsa-miR-324-5p	MIMAT0000761	-2,50	8x10 ⁻³
hsa-miR-140-5p	MIMAT0000431	-2,53	1x10 ⁻²
hsa-miR-450a-5p	MIMAT0001545	-2,72	5x10 ⁻²
hsa-miR-142-3p	MIMAT0000434	-3,35	6x10 ⁻³
hsa-miR-192-5p	MIMAT0000222	-4,39	6x10 ⁻⁴
hsa-miR-128-3p	MIMAT0000424	-11,68	6x10 ⁻⁴
hsa-miR-422a	MIMAT0001339	-12,62	2x10 ⁻³
hsa-miR-143-3p	MIMAT0000435	-20,19	6x10 ⁻⁴
hsa-miR-579-3p	MIMAT0003244	-20,20	8x10 ⁻³
REL-R vs. HD	Accession number	FC	Adjusted P-value
hsa-miR-451a	MIMAT0001631	18,17	1x10 ⁻³
hsa-miR-144-3p	MIMAT0000436	7,97	5x10 ⁻³
hsa-miR-34a-5p	MIMAT0000255	4,16	5x10 ⁻³
hsa-miR-196b-5p	MIMAT0001080	4,22	1x10 ⁻²
hsa-miR-520d-5p/527/518a-5p	MIMAT0002855	-2,45	3x10 ⁻²
hsa-miR-574-5p	MIMAT0004795	-2,74	3x10 ⁻²
hsa-miR-342-3p	MIMAT0000753	-3,27	3x10 ⁻²
hsa-miR-6724-5p	MIMAT0025856	-2,50	4x10 ⁻²
REL-R vs. REL-D	Accession number	FC	Adjusted P-value
hsa-miR-579-3p	MIMAT0003244	3,52	1
hsa-miR-143-3p	MIMAT0000435	2,90	1
hsa-miR-450a-5p	MIMAT0001545	2,30	1
hsa-miR-378e	MIMAT0018927	2,27	1
hsa-miR-145-5p	MIMAT0000437	2,19	1
hsa-miR-4516	MIMAT0019053	1,61	1
hsa-miR-363-3p	MIMAT0000707	1,57	1
hsa-miR-365a/b-3p	MIMAT0000710	1,46	1
hsa-miR-664a-3p	MIMAT0005949	-1,35	1
hsa-let-7a-5p	MIMAT0000062	-1,38	1
hsa-miR-222-3p	MIMAT0000279	-1,48	1
hsa-let-7c-5p	MIMAT0000064	-1,59	1
hsa-miR-181a-5p	MIMAT0000256	-1,60	1
hsa-let-7b-5p	MIMAT0000063	-1,84	1
hsa-miR-100-5p	MIMAT0000098	-1,86	1
hsa-miR-1246	MIMAT0005898	-2,09	1
hsa-miR-9-5p	MIMAT0000441	-2,13	1
hsa-miR-125b-5p	MIMAT0000423	-2,20	1
hsa-miR-10a-5p	MIMAT0000253	-2,53	1

hsa, *homo sapiens*; miR, microRNA; REL-D, patient at the time of diagnosis; REL-R, relative samples at relapse; NREL, patients at diagnosis who never presented relapse; HD, healthy donors.

AML cell lines. It was first analysed whether the combination of JQ1 and curcumin could modulate miR-221-3p and miR-222-3p expression in AML cell lines.

RT-qPCR experiments (Fig. 2A), revealed a trend towards downregulation of the two miRs in all the cell lines, featuring

an enhanced effect with the drug combination. The expression of miR-221-3p expression showed a clear trend towards downregulation, reaching statistical significance in MV-4-11 cells when treated with JQ1, curcumin or the combination of both (Fig. 2A, upper panel). miR-222-3p demonstrated a

Table III. miR expression signatures related to relapse risk or not-relapse with validated targets.

RELAPSE RISK	Validated target genes	NOT RELAPSE	Validated target genes
hsa-miR-128-3p		hsa-miR-200c-3p	ZEB1; ERRFI1
hsa-miR-143-3p	KRAS; FNDC3B; DNMT3A; MAPK7	hsa-miR-191-5p	
hsa-miR-142-3p		hsa-miR-194-5p	
hsa-miR-451a	MIF; ABCB1	hsa-miR-6724-5p	
hsa-miR-29c-3p		hsa-miR-520f-3p	
hsa-miR-140-5p	HDAC4; IGFBP5; VEGFA	hsa-miR-1290	
hsa-miR-196b-5p		hsa-miR-421	
hsa-miR-221-3p	BMF; FOS; CDKN1B; KIT; ESR1; CDKN1C; ICAM1; DDIT4	hsa-miR-365a/b-3p	
hsa-miR-222-3p	CDKN1B; FOS; KIT; CDKN1C; ESR1; MMP1; SOD2; PPP2R2A	hsa-miR-612	
hsa-miR-324-5p	SMO; GLI1	hsa-miR-150-5p	
hsa-miR-34a-5p	MAP2K1; E2F3; SIRT1; MYB; CDK6; DLL1; CCND1; NOTCH1; NOTCH2; JAG1; MET; BCL2; MYCN; WNT1; AXIN2; VEGFA; MYC	hsa-miR-155-5p	JARID2; IKBKE; ETS1; BACH1; TAB2; MEIS1; MECP2; CEBPB; FOXO3; FGF7; SOCS1; INPP5D; AGTR1; SPI1; CYR61; SMAD2; LDOC1; MATR3; TM6SF1; RHOA
		hsa-miR-342-3p	
		hsa-miR-4286	
		hsa-miR-1285-5p	
		hsa-miR-4284	
		hsa-miR-575	
		hsa-miR-1973	

hsa, *homo sapiens*; miR, microRNA.

trend towards downregulation with coupled treatment in all cell-lines, but it did not reach statistical significance (Fig. 2A, lower panel).

It was then analysed whether the combination of JQ1 and curcumin could modulate the CDKN1B protein level, that it is a miR-221-3p and miR-222-3p target, as before mentioned. As revealed in Fig. 2B, the treatment determined an increase in CDKN1B expression both in THP-1 and in MOLM-13 cells while in MV-411 cells the effect of drug combination on the upregulation of CDKN1B expression was comparable to that obtained by single treatment with JQ1.

Association of curcumin and JQ1 causes apoptosis. It was assessed whether these treatments could drive the cell lines toward an apoptotic response. A significant increase in AnnV positive cell percentage was identified in all the cell lines while treated with the drug combination compared with DMSO (Figs. S1 and 3). At a deeper glance, THP-1 and MV-4-11 cells, showing a minor increase in CDKN1B expression, displayed a milder apoptotic response compared with MOLM-13, characterized instead by higher changes in term of expression of CDKN1B (Figs. 2 and 3).

CD34⁺ primary cells from AML patients exhibit miR-221-3p and miR-222-3p downregulation after single and

double treatments. miR-221-3p and miR-222-3p expression was analysed in CD34⁺ cells isolated from BM samples from three randomly selected patients from our cohort. The purity of CD34⁺ cells after isolation was 89% (Fig. S2). After CD34⁺ cells were cultured with MethoCult methylcellulose medium supplemented with JQ1 and curcumin, RNA was extracted and miR-221-3p, miR-222-3p expression was evaluated. In all the samples analysed miR-221-3p and miR-222-3p were modulated when treated with JQ1 and curcumin. In particular, the double treatment led to a significant downregulation of their expression (Fig. 4).

Discussion

Despite improvements in the treatment of paediatric patients affected by AML, novel therapeutic strategies are needed, particularly for paediatric high-risk AML, characterized by high relapse incidence. miRs have been extensively studied as potential biomarkers in adult AML and, recently, expression signatures of miRs in paediatric samples have been proposed (14,19,42).

Notably, few miRNA-based prognostic models have been proposed for paediatric cytogenetically normal AML (18,43), t(8;21) RUNX1-RUNX1T1 AML (44) and AML without considering the cytogenetics (18,45). However, conflicting




Figure 1. NanoString Analysis reveals shared and non-shared miR expression signatures in pediatric patients with AML. (A) Supervised hierarchical clustering of REL-D vs. NREL analysis showed miR expression signature for relapse risk correlation at the time of diagnosis. Only miR with $P \leq 0.05$ were plotted. Benjamini-Hochberg multiple comparisons statistical method was used to control False Discovery Rate. (B) Venn diagrams reporting shared and non-shared dysregulated miRs between REL-D vs. NREL, REL-D vs. HD and NREL vs. HD analysis.

results on the prognostic value of different miRs were reported, possibly due to the vast variability of miRs expression among different cytogenetic and molecular subtypes

of AML (15). It was therefore decided, first, to characterize signature of miRs in patients harbouring *FLT3-ITD* and *MLL* rearrangement to verify whether distinct AML




Figure 2. Curcumin and JQ1 singular and coupled treatments impair protein and miR-221-3p and miR-222-3p expression in leukemia cell lines. (A) Reverse transcription-quantitative PCR showed a significant decrease of miR-221-3p in MV-4-11 cells after 48 h of coupled treatment (ANOVA followed by Tukey's post hoc test), while miR-222-3p was not significantly downregulated. (B) Western blot analysis revealing the expression of CDKN1B at protein level after single and coupled treatment with curcumin and JQ1 in protein lysates of AML cell lines. CDKN1B displayed a significant increase in its expression while treated with the combination drugs and mildly with singular drugs. All the western blots were performed after 48 h from treatment with 250 nM of JQ1 and 10 μ M curcumin. * $P \leq 0.05$, ** $P \leq 0.01$ and *** $P \leq 0.005$.

molecular subsets could affect different miR expression fingerprints. Moreover, to further narrow our miRs signature (identifying only AML associated miRs), HDs were used as a control group. Data on expression of miRs among different AML molecular subtypes are available (16). In accordance with literature data (46), miR-9-5p and miR-10a-5p were upregulated in *MLL*-rearranged and *FLT3-ITD* sets, respectively, even compared with HDs. In addition, miR-99a-5p was upregulated in *FLT3-ITD* patients as compared with *MLL* patients, and was already described in literature as a potential oncomiR in paediatric AML (42). miR-196b-5p and miR-34a-5p were both upregulated in *MLL* and *FLT3-ITD* with respect to HDs and they are possibly involved in the pathogenesis of both AML variants. miR-34a is a widely studied miR as a promising target and it has been considered

as a preclinical and clinical model for the treatment of solid tumours, myeloma and B-cell lymphoma (47,48).

Despite the limited number of patients, to the best of our knowledge, this is the first study investigating the role of miRs as predictive biomarker of relapse in paediatric AML patients with *FLT3-ITD* or *MLL* rearrangement. A total of 28 miRs differentially expressed were identified between patients who relapsed and those who did not, being possibly associated with prognosis. Among them, the vast majority of miRs have already been reported as related to cancer. Comparing the signature of our miRs to those previously reported, a common dysregulation of miR-155 was identified, but displaying an opposite behaviour with a protective effect in the present study as opposed to a relapse association in others (19,49). miR-155 has led to conflicting results in previous studies,




Figure 3. Curcumin and JQ1 treatments lead leukemia cell lines to apoptosis. Annexin V positive cells assessed by flow cytometry were significantly increased in all the cell lines after 48 h coupled treatment with 10 μ M curcumin and 250 nM JQ1 (ANOVA followed by Tukey's post hoc test). * $P \leq 0.05$ and *** $P \leq 0.005$.




Figure 4. Primary CD34⁺ cultures cells exhibit a decrease of miR-221-3p and miR-222-3p expression after treatments. Expression of miR-221-3p and miR-222-3p was significantly downregulated in CD34⁺ primary cells when treated with both JQ1 and curcumin (ANOVA followed by Dunnett's test). * $P \leq 0.05$ and *** $P \leq 0.005$.

with certain groups reporting an anti-leukemic role (50) and others (13,51-53) reporting a role as oncomiR in AML. This is possibly explained by a different level of expression of

miR-155, acting as a tumour suppressor when highly expressed and as an oncomiR when overexpressed to an intermediate level (54). miR-34a-5p expression was positively correlated

with relapse. It was already described that patients with low miR-34a expression showed shorter overall and recurrence-free survival (55). Numerous of the identified miRs, have also been independently considered as promising tool to develop novel therapies; these include miR-200c (47,56), which was revealed to be associated to NREL patients.

miR-221-3p and miR-222-3p, both associated to relapse, gained the attention of the authors as they have been broadly reported in literature as oncomiRs both in hematologic malignancies such as chronic lymphocytic leukemia (57), myelodysplastic syndrome (58), acute lymphoblastic leukemia (59) and AML (60,61) as well as in solid tumours. These miRs have as their primary target CDKN1B, a master regulator of the cell cycle. CDKN1B is a well-known cyclin-dependent kinase inhibitor which regulates cell cycle progression at G1 stage, preventing the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, resulting in a blockade of cell division cycle (62).

BET and HAT inhibitors have already been associated with modulation of miRs in hematological malignancies (63,64). The present findings revealed, for the first time, that JQ1 determines a clear trend towards downregulation of miR-221-3p expression in both MOLM-13 and MV-4-11 AML cell lines with a synergic effect when associated with curcumin; reaching statistical significance in the second one. This is interesting also considering that for *FLT3-ITD* rearrangement MOLM-13 expresses both mutated and wild-type allele, while MV-4-11 expresses mutated allele only (65). It could be hypothesized that *FLT3-ITD* in both mutated alleles renders MV-4-11 cells more sensitive to the effect of drug on miR-221-3p modulation.

Moreover, following the combined treatment, an increase was demonstrated in CDKN1B expression and in apoptotic response in our AML cell lines. The present results were confirmed in cultures with primary leukemic cells, showing a marked reduction of miR-221-3p and miR-222-3p expression. These results supported the idea that BET inhibitors, along with curcumin, could regulate not only coding RNA transcription, but also non-coding RNA such as miRs.

Although the combination of JQ1 and curcumin synergistically reduced miR-221 and miR-222 expression and increased apoptosis in AML cells, a limitation to the present study was represented by insufficient patient samples. Direct regulation of CDKN1B by miR-221-3p and miR-222-3p should be confirmed by further experiments including western blot analysis to verify expression levels of CDKN1B in samples of patients and HDs and the use of inhibitors of miR-221 or miR-222 in a bigger cohort of patients.

In conclusion, the present study identified fingerprints of miRs related to relapse and non-relapse in paediatric *FLT3-ITD*- or *MLL*-rearranged AML. Numerous of these miRs are known to be involved in pathogenetic mechanisms of several haematological malignancies as well as solid tumours and represent both good candidates for targeted treatments and therapeutic tools in different neoplasms. The use of the well-known BRD4 inhibitor JQ1, as well as novel BRD inhibitors in the care of leukaemia could be potentiated by epigenetic drugs such as HATs inhibitors, antagomiR or miR mimic and could expand the therapeutic arsenal in HR-AMLs, particularly for paediatric patients.

Acknowledgements

The authors would like to thank Giuseppe Basso (1948-2021), MD, Department of Woman's and Children's Health, University of Padua (Padua, Italy) for his help in the study design and for providing samples.

Funding

The present study was supported by Ministero della Salute (grant no. GR-2011-02350175) and by Fondazione Umberto Veronesi (Milano, Italy).

Availability of data and materials

The datasets generated and/or analyzed during the current study are available in the GEO repository (accession no. GSE209871; <https://www.ncbi.nlm.nih.gov/gds>).

Authors' contributions

PPL, PV, AB, PM and DP designed the study. RR, AB, PPL and PV conceived and designed the experiments. PPL, PV and VT performed the experiments and data analyses. GN, DV, PV, PF, MS and SR performed Nanostring, data and statistical analyses. VP, FS, EM, AB, DP and PM enrolled patients and collected clinical samples. SR, PV, PPL, MM and PM contributed to manuscript preparation, editing and reviewing. GN and PPL confirm the authenticity of all the raw data. All authors read and approved the final version of the manuscript.

Ethics approval and consent to participate

The present study was approved by the Institutional Review Boards of Bambino Gesù Children's Hospital (Rome, Italy). Informed consent was obtained from either parents or legal guardians according to the Declaration of Helsinki.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Pui CH, Carroll WL, Meshinchi S and Arceci RJ: Biology, risk stratification, and therapy of pediatric acute leukemias: An update. *J Clin Oncol* 29: 551-565, 2011.
2. Creutzig U, van den Heuvel-Eibrink MM, Gibson B, Dworzak MN, Adachi S, de Bont E, Harbott J, Hasle H, Johnston D, Kinoshita A, *et al*: Diagnosis and management of acute myeloid leukemia in children and adolescents: Recommendations from an international expert panel. *Blood* 120: 3187-3205, 2012.
3. Rubnitz JE and Inaba H: Childhood acute myeloid leukaemia. *Br J Haematol* 159: 259-276, 2012.
4. Elgarten CW and Aplenc R: Pediatric acute myeloid leukemia: Updates on biology, risk stratification, and therapy. *Curr Opin Pediatr* 32: 57-66, 2020.

5. Buldini B, Rizzatti F, Masetti R, Fagioli F, Menna G, Micalizzi C, Putti MC, Rizzari C, Santoro N, Zecca M, *et al*: Prognostic significance of flow-cytometry evaluation of minimal residual disease in children with acute myeloid leukaemia treated according to the AIEOP-AML 2002/01 study protocol. *Br J Haematol* 177: 116-126, 2017.
6. Manara E, Basso G, Zampini M, Buldini B, Tregnago C, Rondelli R, Masetti R, Bisio V, Frison M, Polato K, *et al*: Characterization of children with FLT3-ITD acute myeloid leukemia: A report from the AIEOP AML-2002 study group. *Leukemia* 31: 18-25, 2017.
7. Masetti R, Vendemini F, Zama D, Biagi C, Pession A and Locatelli F: Acute myeloid leukemia in infants: Biology and treatment. *Front Pediatr* 3: 37, 2015.
8. Balgobind BV, Raimondi SC, Harbott J, Zimmermann M, Alonzo TA, Auvrignon A, Beverloo HB, Chang M, Creutzig U, Dworzak MN, *et al*: Novel prognostic subgroups in childhood 11q23/MLL-rearranged acute myeloid leukemia: Results of an international retrospective study. *Blood* 114: 2489-2496, 2009.
9. Jiang X, Huang H, Li Z, Li Y, Wang X, Gurbuxani S, Chen P, He C, You D, Zhang S, *et al*: Blockade of miR-150 maturation by MLL-fusion/MYC/LIN-28 is required for MLL-associated leukemia. *Cancer Cell* 22: 524-535, 2012.
10. Ambros V: The functions of animal microRNAs. *Nature* 431: 350-355, 2004.
11. Bartel DP: MicroRNAs: Target recognition and regulatory functions. *Cell* 136: 215-233, 2009.
12. Di Leva G, Garofalo M and Croce CM: MicroRNAs in cancer. *Annu Rev Pathol* 9: 287-314, 2014.
13. Marcucci G, Mrózek K, Radmacher MD, Garzon R and Bloomfield CD: The prognostic and functional role of microRNAs in acute myeloid leukemia. *Blood* 117: 1121-1129, 2011.
14. Obulkasim A, Katsman-Kuipers JE, Verboon L, Sanders M, Touw I, Jongen-Lavrencic M, Pieters R, Klusmann JH, Michel Zwaan C, van den Heuvel-Eibrink MM and Fornerod M: Classification of pediatric acute myeloid leukemia based on miRNA expression profiles. *Oncotarget* 8: 33078-33085, 2017.
15. Danen-van Oorschot AA, Kuipers JE, Arentsen-Peters S, Schotte D, de Haas V, Trka J, Baruchel A, Reinhardt D, Pieters R, Zwaan CM and van den Heuvel-Eibrink MM: Differentially expressed miRNAs in cytogenetic and molecular subtypes of pediatric acute myeloid leukemia. *Pediatr Blood Cancer* 58: 715-721, 2012.
16. Wallace JA and O'Connell RM: MicroRNAs and acute myeloid leukemia: Therapeutic implications and emerging concepts. *Blood* 130: 1290-1301, 2017.
17. Wu D, Qiu Y, Jiao Y, Qiu Z and Liu D: Small molecules targeting HATs, HDACs, and BRDs in cancer therapy. *Front Oncol* 10: 560487, 2020.
18. Zhu R, Zhao W, Fan F, Tang L, Liu J, Luo T, Deng J and Hu Y: A 3-miRNA signature predicts prognosis of pediatric and adolescent cytogenetically normal acute myeloid leukemia. *Oncotarget* 8: 38902-38913, 2017.
19. Lim EL, Trinh DL, Ries RE, Wang J, Gerbing RB, Ma Y, Topham J, Hughes M, Pleasance E, Mungall AJ, *et al*: MicroRNA expression-based model indicates event-free survival in pediatric acute myeloid leukemia. *J Clin Oncol* 35: 3964-3977, 2017.
20. Brondfield S, Umesh S, Corella A, Zuber J, Rappaport AR, Gaillard C, Lowe SW, Goga A and Kogan SC: Direct and indirect targeting of MYC to treat acute myeloid leukemia. *Cancer Chemother Pharmacol* 76: 35-46, 2015.
21. Benetatos L and Vartholomatos G: MicroRNAs mark in the MLL-rearranged leukemia. *Ann Hematol* 92: 1439-1450, 2013.
22. Bretones G, Delgado MD and León J: Myc and cell cycle control. *Biochim Biophys Acta* 1849: 506-516, 2015.
23. Donati B, Lorenzini E and Ciarrocchi A: BRD4 and cancer: Going beyond transcriptional regulation. *Mol Cancer* 17: 164, 2018.
24. Jiang G, Deng W, Liu Y and Wang C: General mechanism of JQ1 in inhibiting various types of cancer. *Mol Med Rep* 21: 1021-1034, 2020.
25. Wu T, Kamikawa YF and Donohoe ME: Brd4's bromodomains mediate histone H3 acetylation and chromatin remodeling in pluripotent cells through P300 and Brg1. *Cell Rep* 25: 1756-1771, 2018.
26. Fiskus W, Sharma S, Qi J, Shah B, Devaraj SG, Leveque C, Portier BP, Iyer S, Bradner JE and Bhalla KN: BET protein antagonist JQ1 is synergistically lethal with FLT3 tyrosine kinase inhibitor (TKI) and overcomes resistance to FLT3-TKI in AML cells expressing FLT-ITD. *Mol Cancer Ther* 13: 2315-2327, 2014.
27. Mio C, Conzatti K, Baldan F, Allegrì L, Sponziello M, Rosignolo F, Russo D, Filetti S and Damante G: BET bromodomain inhibitor JQ1 modulates microRNA expression in thyroid cancer cells. *Oncol Rep* 39: 582-588, 2018.
28. Schick M, Habringer S, Nilsson JA and Keller U: Pathogenesis and therapeutic targeting of aberrant MYC expression in haematological cancers. *Br J Haematol* 179: 724-738, 2017.
29. Chen H, Liu H and Qing G: Targeting oncogenic Myc as a strategy for cancer treatment. *Signal Transduct Target Ther* 3: 5, 2018.
30. Delmore JE, Issa GC, Lemieux ME, Rahl PB, Shi J, Jacobs HM, Kastiris E, Gilpatrick T, Paranal RM, Qi J, *et al*: BET bromodomain inhibition as a therapeutic strategy to target c-Myc. *Cell* 146: 904-917, 2011.
31. Marcu MG, Jung YJ, Lee S, Chung EJ, Lee MJ, Trepel J and Neckers L: Curcumin is an inhibitor of p300 histone acetyltransferase. *Med Chem* 2: 169-174, 2006.
32. Kang SK, Cha SH and Jeon HG: Curcumin-induced histone hypoacetylation enhances caspase-3-dependent glioma cell death and neurogenesis of neural progenitor cells. *Stem Cells Dev* 15: 165-174, 2006.
33. Morimoto T, Sunagawa Y, Kawamura T, Takaya T, Wada H, Nagasawa A, Komeda M, Fujita M, Shimatsu A, Kita T and Hasegawa K: The dietary compound curcumin inhibits p300 histone acetyltransferase activity and prevents heart failure in rats. *J Clin Invest* 118: 868-878, 2008.
34. Spriano F, Stathis A and Bertoni F: Targeting BET bromodomain proteins in cancer: The example of lymphomas. *Pharmacol Ther* 215: 107631, 2020.
35. Wang X, Yang Y, Ren D, Xia Y, He W, Wu Q, Zhang J, Liu M, Du Y, Ren C, *et al*: JQ1, a bromodomain inhibitor, suppresses Th17 effectors by blocking p300-mediated acetylation of ROR γ t. *Br J Pharmacol* 177: 2959-2973, 2020.
36. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
37. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W and Smyth GK: limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* 43: e47, 2015.
38. Reich M, Liefeld T, Gould J, Lerner J, Tamayo P and Mesirov JP: GenePattern 2.0. *Nat Genet* 38: 500-501, 2006.
39. Heberle H, Meirelles GV, da Silva FR, Telles GP and Minghim R: InteractiVenn: A web-based tool for the analysis of sets through Venn diagrams. *BMC Bioinformatics* 16: 169, 2015.
40. Dweep H and Gretz N: miRWalk2.0: A comprehensive atlas of microRNA-target interactions. *Nat Methods* 12: 697, 2015.
41. Haferlach C, Kern W, Schindela S, Kohlmann A, Alpermann T, Schnittger S and Haferlach T: Gene expression of BAALC, CDKN1B, ERG, and MN1 adds independent prognostic information to cytogenetics and molecular mutations in adult acute myeloid leukemia. *Genes Chromosomes Cancer* 51: 257-265, 2012.
42. Zhang H, Luo XQ, Zhang P, Huang LB, Zheng YS, Wu J, Zhou H, Qu LH, Xu L and Chen YQ: MicroRNA patterns associated with clinical prognostic parameters and CNS relapse prediction in pediatric acute leukemia. *PLoS One* 4: e7826, 2009.
43. Gaur V, Chaudhary S, Tyagi A, Agarwal S, Sharawat SK, Sarkar S, Singh H, Bakhshi S, Sharma P and Kumar S: Dysregulation of miRNA expression and their prognostic significance in paediatric cytogenetically normal acute myeloid leukaemia. *Br J Haematol* 188: e90-e94, 2020.
44. Zampini M, Bisio V, Leszl A, Putti MC, Menna G, Rizzari C, Pession A, Locatelli F, Basso G, Tregnago C and Pigazzi M: A three-miRNA-based expression signature at diagnosis can predict occurrence of relapse in children with t(8;21) RUNX1-RUNX1T1 acute myeloid leukaemia. *Br J Haematol* 183: 298-301, 2018.
45. Zhu R, Lin W, Zhao W, Fan F, Tang L and Hu Y: A 4-microRNA signature for survival prognosis in pediatric and adolescent acute myeloid leukemia. *J Cell Biochem* 120: 3958-3968, 2019.
46. Liu Y, Cheng Z, Pang Y, Cui L, Qian T, Quan L, Zhao H, Shi J, Ke X and Fu L: Role of microRNAs, circRNAs and long noncoding RNAs in acute myeloid leukemia. *J Hematol Oncol* 12: 51, 2019.
47. Rupaimoole R and Slack FJ: MicroRNA therapeutics: Towards a new era for the management of cancer and other diseases. *Nat Rev Drug Discov* 16: 203-222, 2017.
48. Beg MS, Brenner AJ, Sachdev J, Borad M, Kang YK, Stoudemire J, Smith S, Bader AG, Kim S and Hong DS: Phase I study of MRX34, a liposomal miR-34a mimic, administered twice weekly in patients with advanced solid tumors. *Invest New Drugs* 35: 180-188, 2017.

49. Yan W, Xu L, Sun Z, Lin Y, Zhang W, Chen J, Hu S and Shen B: MicroRNA biomarker identification for pediatric acute myeloid leukemia based on a novel bioinformatics model. *Oncotarget* 6: 26424-26436, 2015.
50. Palma CA, Al Sheikha D, Lim TK, Bryant A, Vu TT, Jayaswal V and Ma DD: MicroRNA-155 as an inducer of apoptosis and cell differentiation in acute myeloid leukaemia. *Mol Cancer* 13: 79, 2014.
51. Marcucci G, Maharry KS, Metzeler KH, Volinia S, Wu YZ, Mrózek K, Nicolet D, Kohlschmidt J, Whitman SP, Mendler JH, *et al*: Clinical role of microRNAs in cytogenetically normal acute myeloid leukemia: miR-155 upregulation independently identifies high-risk patients. *J Clin Oncol* 31: 2086-2093, 2013.
52. Wallace JA, Kagele DA, Eiring AM, Kim CN, Hu R, Runtsch MC, Alexander M, Huffaker TB, Lee SH, Patel AB, *et al*: miR-155 promotes FLT3-ITD-induced myeloproliferative disease through inhibition of the interferon response. *Blood* 129: 3074-3086, 2017.
53. O'Connell RM, Rao DS, Chaudhuri AA, Boldin MP, Taganov KD, Nicoll J, Paquette RL and Baltimore D: Sustained expression of microRNA-155 in hematopoietic stem cells causes a myeloproliferative disorder. *J Exp Med* 205: 585-594, 2008.
54. Narayan N, Morenos L, Phipson B, Willis SN, Brumatti G, Eggers S, Lalaoui N, Brown LM, Kosasih HJ, Bartolo RC, *et al*: Functionally distinct roles for different miR-155 expression levels through contrasting effects on gene expression, in acute myeloid leukaemia. *Leukemia* 31: 808-820, 2017.
55. Huang Y, Zou Y, Lin L, Ma X and Chen H: Identification of serum miR-34a as a potential biomarker in acute myeloid leukemia. *Cancer Biomark* 22: 799-805, 2018.
56. Chakraborty C, Sharma AR, Sharma G, Doss CGP and Lee SS: Therapeutic miRNA and siRNA: Moving from bench to clinic as next generation medicine. *Mol Ther Nucleic Acids* 8: 132-143, 2017.
57. Frenquelli M, Muzio M, Scielzo C, Fazi C, Scarfò L, Rossi C, Ferrari G, Ghia P and Caligaris-Cappio F: MicroRNA and proliferation control in chronic lymphocytic leukemia: Functional relationship between miR-221/222 cluster and p27. *Blood* 115: 3949-3959, 2010.
58. Hussein K, Theophile K, Büsche G, Schlegelberger B, Göhring G, Kreipe H and Bock O: Significant inverse correlation of microRNA-150/MYB and microRNA-222/p27 in myelodysplastic syndrome. *Leuk Res* 34: 328-334, 2010.
59. Moses BS, Evans R, Slone WL, Piktet D, Martinez I, Craig MD and Gibson LF: Bone marrow microenvironment niche regulates miR-221/222 in acute lymphoblastic leukemia. *Mol Cancer Res* 14: 909-919, 2016.
60. Rommer A, Steinleitner K, Hackl H, Schneckenleithner C, Engelmann M, Scheideler M, Vlatkovic I, Kralovics R, Cerny-Reiterer S, Valent P, *et al*: Overexpression of primary microRNA 221/222 in acute myeloid leukemia. *BMC Cancer* 13: 364, 2013.
61. Lee YG, Kim I, Oh S, Shin DY, Koh Y and Lee KW: Small RNA sequencing profiles of mir-181 and mir-221, the most relevant microRNAs in acute myeloid leukemia. *Korean J Intern Med* 34: 178-183, 2019.
62. le Sage C, Nagel R and Agami R: Diverse ways to control p27Kip1 function: miRNAs come into play. *Cell Cycle* 6: 2742-2749, 2007.
63. Kohnken R, McNeil B, Wen J, McConnell K, Grinshpun L, Keiter A, Chen L, William B, Porcu P and Mishra A: Preclinical targeting of MicroRNA-214 in cutaneous T-cell lymphoma. *J Invest Dermatol* 139: 1966-1974.e3, 2019.
64. Mensah AA, Cascione L, Gaudio E, Tarantelli C, Bomben R, Bernasconi E, Zito D, Lampis A, Hahne JC, Rinaldi A, *et al*: Bromodomain and extra-terminal domain inhibition modulates the expression of pathologically relevant microRNAs in diffuse large B-cell lymphoma. *Haematologica* 103: 2049-2058, 2018.
65. Quentmeier H, Reinhardt J, Zaborski M and Drexler HG: FLT3 mutations in acute myeloid leukemia cell lines. *Leukemia* 17: 120-124, 2003.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.