

Positive immunohistochemical staining of γ H2AX is associated with tumor progression in gastric cancers from radiation-exposed patients

KAZUHIRO SENTANI¹, NAOHIDE OUE¹, NAOYA SAKAMOTO¹, TAKASHI NISHISAKA²,
TOSHIYUKI FUKUHARA², HIROO MATSUURA³ and WATARU YASUI¹

¹Department of Molecular Pathology, Hiroshima University Graduate School of Biomedical Sciences, Hiroshima;

²Department of Pathology and Laboratory Medicine, Hiroshima Prefectural Hospital, Hiroshima;

³Department of Pathology, Hiroshima City Hospital, Hiroshima, Japan

Received May 27, 2008; Accepted August 2, 2008

DOI: 10.3892/or_00000120

Abstract. To elucidate the mechanism of radiation-induced cancers, molecular analysis of cancers in atomic bomb (A-bomb) exposure is important. DNA double-strand breaks (DSBs) are thought to be caused by the deleterious effects of ionizing radiation, and γ H2AX (serine 139 phosphorylated form of histone H2AX) is reported to be a significant marker for DSBs. In the present study, we performed immunohistochemical analysis of γ H2AX in gastric cancers (GCs) from 66 exposed and 47 non-exposed patients who developed GC after the bombing. Of the 47 GCs from non-exposed patients, 6 (13%) cases showed nuclear positive staining for γ H2AX, whereas of the 66 GCs from exposed patients, 20 (30%) cases were positive ($P=0.0405$). However, among stage I GC, there was no significant difference in γ H2AX expression frequency between exposed patients and non-exposed patients. Among exposed patients, stage II-IV cases were more frequently positive for γ H2AX than stage I cases ($P=0.0197$). Among GCs from non-exposed patients, γ H2AX staining showed no significant association with Lauren's classification, depth of invasion, lymph node metastasis or TNM stage. These results suggest that the characteristics of tumor cells differ between GCs from exposed and non-exposed patients. DSBs may be involved in progression of GC in exposed patients.

Introduction

More than 60 years have passed since atomic bomb (A-bombs) exposure in Hiroshima and Nagasaki, Japan. A

prospective cohort study (Life Span Study, LSS) of 120,000 subjects is being conducted by the Radiation Effects Research Foundation (RERF) (1). It was reported that exposure to ionizing radiation (IR) increases the risk of leukemia and other cancers (2), and damage to nuclear DNA likely represents an initiating event for carcinogenesis. Increases in cancer risk due to exposure to IR are based on epidemiologic studies of exposed human populations, mainly the A-bomb survivors of Hiroshima and Nagasaki (3). Solid cancers, including breast, colon, lung and stomach cancers, have a long latency period, and the excess relative risks (RRs) of solid cancers remain high, specifically among those exposed when young (1). Although approximately half of the LSS members are now deceased, cancer mortality in the LSS has continued to increase as this population ages, and it is anticipated to peak in 2015.

According to the World Health Organization, gastric cancer (GC) is the fourth most common malignancy worldwide, with approximately 870,000 new cases occurring yearly. Cancer develops as a result of multiple genetic and epigenetic alterations (4,5). Although several genetic alterations, including mutations in *TP53* and *BRAF*, have been reported in selected cancers of A-bomb survivors (6-8), specific mutations for radiation-associated cancers have not been reported.

DNA double-strand breaks (DSBs) are thought to be caused by the deleterious effects of IR (9,10). DSBs can induce chromosomal aberrations that cause cells to malfunction, resulting in cell death or tumorigenesis (10). One of the earliest steps in the cellular response to DSBs is the phosphorylation of histone H2AX at serine 139 (γ H2AX), the site of γ -phosphorylation (11). H2AX can be phosphorylated by several phosphoinositide-3 (PI3) kinases including ataxia telangiectasia mutated (ATM), DNA-dependent protein kinase (DNA-PK) and ataxia telangiectasia and Rad3 related (ATR) (12). The number of resulting γ H2AX foci has been correlated directly with the number of DSBs produced by IR (13,14). Therefore, the number of γ H2AX foci is a significant marker for DSBs. Immunohistochemical analyses of γ H2AX have been reported for human cancers of the urinary bladder,

Correspondence to: Dr Wataru Yasui, Department of Molecular Pathology, Hiroshima University Graduate School of Biomedical Sciences, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8551, Japan
E-mail: wyasui@hiroshima-u.ac.jp

Key words: γ H2AX, gastric cancer, radiation, atomic bomb, DNA double-strand break

breast, lung, colon and prostate (15-17). It was also reported that γ H2AX-positive cells are present in colorectal cancer (CRC) and precursor lesions, such as adenoma, but not in normal colonic epithelium (15). Furthermore, invasive CRCs were reported to show less γ H2AX staining than adenomas (15). These results suggest that staining of γ H2AX correlates with DNA damage checkpoint activation in premalignant lesions. Therefore, the existence of γ H2AX foci might be a useful and sensitive marker of cancer, especially for detecting cancers or precursor lesions in A-bomb survivor, because IR induces DSBs. However, there are no reports of immunohistochemical analyses of γ H2AX in GCs from either IR-exposed patients or -non-exposed patients. Therefore, in the present study, we performed immunohistochemical analysis of γ H2AX in 113 GCs derived from A-bomb survivors.

Patients and methods

Patients and tumor specimens. For immunohistochemical analysis, we used formalin-fixed, paraffin-embedded archival tissues from 113 patients with GC who underwent surgery between 1975 and 2005 at Hiroshima University Hospital (Hiroshima, Japan) or an affiliated hospital. Only patients who did not undergo preoperative radio- or chemotherapy were enrolled in the study. All 113 patients were A-bomb survivors (LSS cohort members) in Hiroshima, Japan. Although these patients were survivors who developed GC after the bombing, they were further classified according to their level of radiation exposure (i.e., ≥ 5 mGy and < 5 mGy were defined as 'exposed' and 'non-exposed', respectively). Our population comprised 66 exposed (median dose, 51 mGy; range, 5-2601 mGy) and 47 non-exposed patients (median dose, 0 mGy; range, 0-4 mGy).

Tumor staging was performed according to the Union Internationale Contre le Cancer (UICC) system (18). Histologic classification was carried out according to the Lauren classification system (19). The detailed procedures for acquiring informed consent from study patients and collecting tissue specimens were described previously (20). In accordance with the Ethical Guidelines For Human Genome/ Gene Research enacted by the Japanese Government, tissue specimens were collected and used after approval from the Ethical Review Committee of the Hiroshima University School of Medicine and from the ethical review committees of collaborating organizations.

Radiation dose. A-bomb radiation doses were estimated with the DS02 system (21).

Immunohistochemistry. From each patient, one representative tumor block, including the tumor center and invasive front as well as tumor-associated non-neoplastic mucosa, was examined by immunohistochemistry. In cases of large, late-stage tumors, different sections were examined to include representative areas of the tumor center as well as of the lateral and deep tumor invasive fronts.

Immunohistochemical detection of γ H2AX was performed with a mouse monoclonal antibody (Upstate Biotechnology, Chicago, IL, USA) and Dako Envision Kit (Dako, Carpinteria, CA). In brief, sections were pretreated by microwaving

(500 W) in citrate buffer (pH 6.0) for 15 min to retrieve antigenicity. After endogenous peroxidase activity was blocked with 3% H₂O₂-methanol for 10 min, sections were incubated with normal goat serum (Dako) for 20 min to block non-specific antibody binding sites. Sections were then incubated with anti- γ H2AX (diluted 1:200) for 1 h at room temperature followed by incubation with peroxidase-labelled anti-mouse IgG for 60 min. Staining was completed with a 10-min incubation with the substrate-chromogen solution. Sections were counterstained with 0.1% hematoxylin. Appropriate negative controls were created by omission of the primary antibody. All slices were evaluated without knowledge of the clinical data.

Double immunofluorescence staining. Double immunofluorescence staining for dewaxed sections was performed with mouse monoclonal anti- γ H2AX antibody (Upstate) with rabbit polyclonal anti-H2AX antibody (Upstate) or mouse monoclonal anti- γ H2AX antibody with a rabbit polyclonal antibody against the activated form of caspase-3 (Promega, Madison, WI, USA). Microwave pretreatment in citrate buffer was performed for 15 min to retrieve antigenicity. Sections were then incubated with normal goat serum for 30 min to block non-specific antibody binding sites. Sections were treated consecutively at room temperature with primary antibodies for 60 min, and immunocomplexes were detected with Alexa Fluor 488-conjugated goat anti-mouse IgG and Alexa Fluor 546-conjugated goat anti-rabbit IgG (Molecular Probes, Eugene, OR, USA).

Statistical methods. Associations between clinicopathologic variables and immunostaining for γ H2AX were analyzed by Fisher's exact test. A P-value < 0.05 was considered statistically significant.

Results

Of 113 GC from A-bomb survivors, 48 (42%) showed nuclear staining of γ H2AX (Fig. 1A). These 48 cases comprised 26 GC cases with diffuse staining for γ H2AX and 22 GC cases with staining of γ H2AX only in superficial portions (Fig. 1B) or in necrotic debris in the lumen (Fig. 1C). We confirmed that γ H2AX yielded granular, nuclear staining (Fig. 1D). H2AX showed ubiquitous staining in GC (Fig. 1E). It was reported previously that γ H2AX is expressed during early apoptosis triggered through the caspase-3/caspase-activated DNase (CAD) pathway (22,23). Double immunofluorescence staining revealed that γ H2AX-positive tumor cells in superficial portions or necrotic debris were also positive for the activated form of caspase-3 (a marker of apoptosis) (Fig. 1F). Because we believed that γ H2AX staining induced by apoptosis was not related to IR, cases with superficial staining and staining of necrotic debris were excluded from the positive cases. In contrast, the percentage of γ H2AX-stained tumor cells was $> 5\%$ in 26 GC cases showing diffuse staining; we considered these as positive cases. Twenty-four of 26 GC cases had from 5% to 10% of γ H2AX-stained tumor cells. In particular, remaining two cases had $> 30\%$ of γ H2AX-stained tumor cells, both of which were α -fetoprotein (AFP)-positive GC.




Figure 1. Immunohistochemical analysis of γ H2AX in gastric carcinoma (GC) tissues (A-F) and non-neoplastic gastric mucosa (G-I). Staining for γ H2AX in GC. Schematic representation of a positive case (A). Tumor cells or necrotic debris with γ H2AX staining are located at the superficial portion of GCs (B) or in the lumen of the tumor gland (C). Granular staining of γ H2AX is found in the nucleus (D). Staining for H2AX. H2AX shows ubiquitous staining both in GC (E) and non-neoplastic gastric mucosa (I). Some surface cells in non-neoplastic gastric mucosa also express γ H2AX (G). Double immunofluorescence staining of the activated form of caspase-3 and γ H2AX. Necrotic debris in the lumen of GCs (F) and superficial apoptotic cells in non-neoplastic gastric mucosa (H) are shown. Cells were imaged with a fluorescence microscope as described in Materials and methods.

We analyzed the association between γ H2AX staining and clinicopathologic parameters in GCs from 66 IR exposed and 47 non-exposed patients (Table I). When all tumor stages were considered, γ H2AX staining was detected in 20 (30%) of 66 exposed patients and 6 (13%) of 47 non-exposed patients ($P=0.0405$). Because IR is a carcinogen and can increase an individual's risk of tumor development, we analyzed immunohistochemical staining for γ H2AX in early-stage GCs. In GCs showing T1 (tumor invades lamina propria or submucosa), N0 (no regional lymph node metastasis), or stage I, γ H2AX positivity did not differ significantly between exposed and non-exposed patients. In contrast,

among T2-4 GCs, γ H2AX was expressed more often in exposed patients than in non-exposed patients ($P=0.0236$). In stage II-IV GCs, γ H2AX expression showed a marginally significant difference between exposed patients and non-exposed patients ($P=0.0602$).

Of the 66 GCs from exposed patients, γ H2AX was present in 20 (30%). γ H2AX expression in GCs from exposed patients was associated significantly with depth of invasion ($P=0.003$) (Table II). Furthermore, γ H2AX staining was observed more frequently in stage II-IV GCs than in stage I GCs ($P=0.0197$) (Table II). In contrast, the presence of γ H2AX in GCs from non-exposed patients showed no signi-

Table I. Expression of γ H2AX in GC and its association with clinicopathologic variables.

	No. of cases	γ H2AX positive (%)	P-value
GC	113	26 (23)	
Exposed patients	66	20 (30)	
Non-exposed patients	47	6 (13)	0.0405
Intestinal type ^a			
Exposed patients	34	9 (26)	
Non-exposed patients	33	5 (15)	0.3689
Diffuse type ^a			
Exposed patients	32	11 (34)	
Non-exposed patients	14	1 (7)	0.073
Depth of invasion ^b			
T1			
Exposed patients	15	0 (0)	
Non-exposed patients	19	2 (11)	0.492
T2-4			
Exposed patients	51	20 (39)	
Non-exposed patients	28	4 (14)	0.0236
Lymph node metastasis ^b			
N0			
Exposed patients	25	5 (20)	
Non-exposed patients	24	2 (8)	0.4174
N1-4			
Exposed patients	41	15 (37)	
Non-exposed patients	23	4 (17)	0.1552
TNM stage ^b			
I			
Exposed patients	21	2 (10)	
Non-exposed patients	23	2 (9)	1.000
II-IV			
Exposed patients	45	18 (40)	
Non-exposed patients	24	4 (17)	0.0602

GC, gastric carcinoma. ^aHistologic classification of GC was according to the Lauren classification system. ^bTumor stage was according to the Tumor-Node-Metastasis (TNM) staging system.

ficant correlation with Lauren's classification, depth of invasion, lymph node metastasis or TNM stage (Table II). There was no significant association between γ H2AX staining and radiation dose at the time of A-bombing (data not shown).

In non-neoplastic gastric mucosa or intestinal metaplasia adjacent to the tumor, only a few superficial cells in both exposed and non-exposed patients showed immunostaining of γ H2AX (Fig. 1G) and activated form of caspase-3 (Fig. 1H). H2AX showed ubiquitous immunostaining (Fig. 1I).

Discussion

While the DNA damage response plays a major role in tumor suppression, how this response contributes to suppression

of stomach tumorigenesis remains unclear. DSBs of chromosomal DNA are thought to be caused by the hazardous effects of IR and may result in chromosomal translocations, deletions or loss of genetic information, which are all causatively linked to tumorigenesis (15). Therefore, DSBs may be involved in radiation-associated gastric carcinogenesis among A-bomb survivors. In the present study, we provide immunohistochemical evidence that γ H2AX is expressed in 13% of GCs from non-exposed patients and 30% of GCs from exposed patients. Because IR is a carcinogen and can increase an individual's risk of tumor development, DSBs appear to play a more important role in early-stage GC rather than late-stage GC. In fact, γ H2AX is reported to be expressed commonly in early precursor lesions in urinary bladder, breast, lung, colon and prostate (15,17). However, in



	No. of cases	γ H2AX positive (%)	P-value
Exposed GC patients	66	20 (30)	
Histology ^a			
Intestinal type	34	9 (26)	
Diffuse type	32	11 (34)	0.5946
Depth of invasion ^b			
T1	15	0 (0)	
T2-4	51	20 (39)	0.003
Lymph node metastasis ^b			
N0	25	5 (20)	
N1-4	41	15 (37)	0.1792
TNM stage ^b			
I	21	2 (10)	
II-IV	45	18 (40)	0.0197
Non-exposed GC patients	47	6 (13)	
Histology ^a			
Intestinal type	33	5 (15)	
Diffuse type	14	1 (7)	0.6532
Depth of invasion ^b			
T1	19	2 (11)	
T2-4	28	4 (14)	1.000
Lymph node metastasis ^b			
N0	24	2 (8)	
N1-4	23	4 (17)	0.4158
TNM stage ^b			
I	23	2 (9)	
II-IV	24	4 (17)	0.6662

GC, gastric carcinoma. ^aHistologic classification of GC was according to the Lauren classification system. ^bTumor stage was done according to be the Tumor-Node-Metastasis (TNM) staging system.

the present study, there was no significant difference in γ H2AX staining between stage I GCs from exposed patients and non-exposed patients. Furthermore, in intestinal metaplasia adjacent to the tumor, which is considered to be a gastric precancerous lesion, staining of γ H2AX was not observed in epithelial or stromal cells. These results suggest that DSBs are less likely to be involved in the genesis of GCs.

In contrast, in exposed patients, γ H2AX-positive GC cases showed more advanced depth of invasion and higher TNM stage than γ H2AX-negative GC cases, suggesting that DSBs may participate in progression of GC in exposed patients. It has been reported that deregulated c-myc expression induces DNA damage and the formation of γ H2AX (24). However, mammalian SWI/SNF complexes facilitate DSBs repair by promoting γ H2AX product (25), and we reported previously that increased expression of BRG1, a component of the SWI/SNF complex, is associated

with advanced-stage GCs (26). It is possible that such signals may also contribute to γ H2AX activation in GCs from exposed patients. Taken together, the molecular mechanisms that underlie phosphorylation of γ H2AX may differ between IR-exposed patients and -non-exposed patients. It is also possible that because a single DSB can result in chromosomal translocations, deletions or loss of genetic information, several genes associated with tumor progression may be deleted in γ H2AX-positive GC cases. Further studies are needed to identify these mechanisms.

In conclusion, DSBs do not appear to be characteristic alterations in stomach carcinogenesis in IR-exposed patients. However, immunohistochemical staining of γ H2AX is increased with tumor progression in GCs from exposed patients. Although it is unclear whether all GCs from exposed patients in the present study were radiation-induced cancers, DSBs may serve as a marker for progression of GCs.

Acknowledgements

We thank Ms. Emiko Hisamoto for excellent technical assistance and advice. This work was carried out with the kind cooperation of the Research Center for Molecular Medicine, Faculty of Medicine, Hiroshima University. We thank the Analysis Center of Life Science, Hiroshima University for the use of their facilities. This work was supported, in part, by Grants-in-Aid for Cancer Research from the Ministry of Education, Culture, Science, Sports, and Technology of Japan; in part by a Grant-in-Aid for the Third Comprehensive 10-Year Strategy for Cancer Control and for Cancer Research from the Ministry of Health, Labour and Welfare of Japan.

References

1. Preston DL, Ron E, Tokuoka S, *et al*: Solid cancer incidence in atomic bomb survivors: 1958-1998. *Radiat Res* 168: 1-64, 2007.
2. Yiin JH, Schubauer-Berigan MK, Silver SR, *et al*: Risk of lung cancer and leukemia from exposure to ionizing radiation and potential confounders among workers at the Portsmouth Naval Shipyard. *Radiat Res* 163: 603-613, 2005.
3. Ron E, Preston DL, Mabuchi K, Thompson DE and Soda M: Cancer incidence in atomic bomb survivors. Part IV: comparison of cancer incidence and mortality. *Radiat Res* 137: 98-112, 1994.
4. Yasui W, Oue N, Kitadai Y and Nakayama H: Recent advances in molecular pathobiology of gastric carcinoma. In: *The Diversity of Gastric Carcinoma Pathogenesis: Diagnosis and Therapy*. Kaminishi M, Takubo K and Mafune K (eds). Springer, Tokyo, pp51-71, 2005.
5. Ushijima T and Sasako M: Focus on gastric cancer. *Cancer Cell* 5: 121-125, 2004.
6. Takeshima Y, Seyama T, Bennett WP, *et al*: p53 mutations in lung cancers from non-smoking atomic-bomb survivors. *Lancet* 342: 1520-1521, 1993.
7. Takahashi K, Eguchi H, Arihiro K, *et al*: The presence of BRAF point mutation in adult papillary thyroid carcinomas from atomic bomb survivors correlates with radiation dose. *Mol Carcinog* 46: 242-248, 2007.
8. Iwamoto KS, Mizuno T, Tokuoka S, Mabuchi K and Seyama T: Frequency of p53 mutations in hepatocellular carcinomas from atomic bomb survivors. *J Natl Cancer Inst* 90: 1167-1168, 1998.
9. Hoeijmakers JH: Genome maintenance mechanisms for preventing cancer. *Nature* 411: 366-374, 2001.
10. Van Gent DC, Hoeijmakers JH and Kanaar R: Chromosomal stability and the DNA double-stranded break connection. *Nat Rev Genet* 2: 196-206, 2001.
11. Rogakou EP, Pilch DR, Orr AH, Ivanova VS and Bonner WM: DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *J Biol Chem* 273: 5858-5868, 1998.
12. Paull TT, Rogakou EP, Yamazaki V, Kirchgessner CU, Gellert M and Bonner WM: A critical role for histone H2AX in recruitment of repair factors to nuclear foci after DNA damage. *Curr Biol* 10: 886-895, 2000.
13. Rothkamm K and Lobrich M: Evidence for a lack of DNA double-strand break repair in human cells exposed to very low x-ray doses. *Proc Natl Acad Sci USA* 100: 5057-5062, 2003.
14. Sedelnikova OA, Rogakou EP, Panyutin IG and Bonner WM: Quantitative detection of (125)IdU-induced DNA double-strand breaks with gamma-H2AX antibody. *Radiat Res* 158: 486-492, 2002.
15. Bartkova J, Horejsi Z, Koed K, *et al*: DNA damage response as a candidate anti-cancer barrier in early human tumorigenesis. *Nature* 434: 864-870, 2005.
16. Gorgoulis VG, Vassiliou LV, Karakaidos P, *et al*: Activation of the DNA damage checkpoint and genomic instability in human precancerous lesions. *Nature* 434: 907-913, 2005.
17. Fan C, Quan R, Feng X, *et al*: ATM activation is accompanied with earlier stages of prostate tumorigenesis. *Biochim Biophys Acta* 1763: 1090-1097, 2006.
18. Sobin LH, Wittekind CH (eds): *TNM Classification of Malignant Tumors*. 6th edition. John Wiley & Sons, New York, pp65-68, 2002.
19. Lauren P: The two histological main types of gastric carcinoma: diffuse and so-called intestinal-type carcinoma. An attempt at a histo-clinical classification. *Acta Pathol Microbiol Scand* 64: 31-49, 1965.
20. Yasui W and Oue N: Systematic collection of tissue specimens and molecular pathological analysis of newly diagnosed solid cancers among atomic bomb survivors. *Int Congr Ser*: 81-86, 2007.
21. Preston DL, Pierce DA, Shimizu Y, *et al*: Effect of recent changes in atomic bomb survivor dosimetry on cancer mortality risk estimates. *Radiat Res* 162: 377-389, 2004.
22. Rogakou EP, Boon C, Redon C and Bonner WM: Megabase chromatin domains involved in DNA double-strand breaks *in vivo*. *J Cell Biol* 146: 905-916, 1999.
23. Lu C, Zhu F, Cho YY, *et al*: Cell apoptosis: requirement of H2AX in DNA ladder formation, but not for the activation of caspase-3. *Mol Cell* 23: 121-132, 2006.
24. Pusapati RV, Rounbehler RJ, Hong S, *et al*: ATM promotes apoptosis and suppresses tumorigenesis in response to Myc. *Proc Natl Acad Sci USA* 103: 1446-1451, 2006.
25. Park JH, Park EJ, Lee HS, *et al*: Mammalian SWI/SNF complexes facilitate DNA double-strand break repair by promoting gamma-H2AX induction. *EMBO J* 25: 3986-3997, 2006.
26. Sentani K, Oue N, Kondo H, *et al*: Increased expression but not genetic alteration of BRG1, a component of the SWI/SNF complex, is associated with the advanced stage of human gastric carcinomas. *Pathobiology* 69: 315-320, 2001.