

Calphostin C, a remarkable multimodal photodynamic killer of neoplastic cells by selective nuclear lamin B1 destruction and apoptosis (Review)

ANNA CHIARINI¹, JAMES F. WHITFIELD², RAFFAELLA PACCHIANA¹,
MADDALENA MARCONI¹, UBALDO ARMATO¹ and ILARIA DAL PRÀ¹

¹Department of Biomedical and Surgical Sciences, University of Verona Medical School, 8 Strada Le Grazie, I-37134 Verona, Italy; ²Institute for Biological Sciences, NRC, 52 Montreal Road, Ottawa, Ontario, K1A-0N6 Canada

Received October 13, 2009; Accepted December 2, 2009

DOI: 10.3892/or_00000711

Abstract. Perylenequinones that generate reactive oxygen species (ROS) when illuminated with visible light have been recommended as photodynamic chemotherapeutic agents. One of these is calphostin C (CalC), the action of the photo-activated derivative of which, CalC^{PE}, has been ascribed to its ability to selectively and irreversibly inhibit protein kinase Cs (PKCs). But recent results of experiments with neoplastic rat fibroblasts and human breast and uterine cervix cancer cells have revealed that the action of CalC^{PE} involves more than PKC inhibition. Besides suppressing PKC activity, CalC^{PE} rapidly causes endoplasmic reticulum (ER) stress in breast cancer cells and the selective complete oxidation and proteasomal destruction of the functionally essential nuclear envelope protein lamin B1, in human cervical carcinoma (HCC) cells and neoplastic rat fibroblasts. When these lamin B1-lacking cells are placed in the dark, cytoplasmic membrane-linked PKC activities suddenly rebound and apoptosis is initiated as indicated by the immediate release of cytochrome *c* from mitochondria and later on the activation of caspases. Hence, CalC^{PE} is a photodynamic cytotoxic agent attacking multiple targets in cancer cells and it would be worth determining, even for their best applicative

use, whether other perylenequinones also share the so far unexpectedly complex deadly properties of the CalC^{PE}.

Contents

1. Introduction
2. The light-induced events and their in-dark sequelae
3. The drivers of CalC^{PE} cytotoxic actions
4. Conclusions

1. Introduction

Perylenequinones are second-generation cell photosensitizers which are easily purified, highly soluble, amenable to site-directed chemical modifications, quickly inactivated in the dark and rapidly cleared from tissues without a persisting tissue photosensitivity. Therefore, the perylenequinones have been recommended as photodynamic therapeutic agents (1-6).

Calphostin C (CalC; UCN-1028c) is one of these agents. It is a dark red to brown, highly lipophilic, protein kinase C (PKC)-inhibiting, perylenequinone metabolite from the fungus *Cladosporium cladosporioides* (Fig. 1A) (7-12). It has a high molecular extinction coefficient (ϵ), a wide UV-visible absorption spectrum with peaks at 226 (ϵ 44,800), 269 (ϵ 29,550), 474 (ϵ 23,550), 553.9 (ϵ shoulder 11,900) and 582 nm (ϵ 12,000), and a high quantum yield of ¹O₂ generation in aqueous solution (5,12,13). According to these properties CalC absorbs enough near infrared light to be photoexcited to CalC^{PE} for use for photodynamic tumor therapy (5).

CalC^{PE} is best known and used for its photodependent strong ability to selectively inhibit most PKC isoforms (7-13). However, it has recently been shown that CalC^{PE} mounts a multi-target attack on ER (endoplasmic reticulum) glycoprotein transport resulting in ER vacuolation, ER stress, and a cluster of apoptogenic consequences [i.e., activation of c-Jun N-terminal kinase and protein kinase R-like ER kinase and upregulation of CCAAT/enhancer-binding protein homologous transcription factor (CHOP/GADD153)] independently of the inhibition of PKCs activity (4). But CalC^{PE} also rapidly and selectively attacks the neoplastic cell nucleus (14).

Correspondence to: Professor Ubaldo Armato, Department of Biomedical and Surgical Sciences, University of Verona Medical School, 8 Strada Le Grazie, I-37134 Verona, Italy
E-mail: ubaldo.armato@univr.it

Abbreviations: ANT, adenine nucleotide translocase; CalC, calphostin C; CalC^{PE}, photoactivated CalC; CPF, cytoplasmic particulate fraction; DAG, diacylglycerol; ER, endoplasmic reticulum; HCC, human cervical carcinoma; NL, nuclear lamina; NMF, nuclear membrane fraction; PKC, protein kinase C; ROS, reactive oxygen species; VDAC, voltage-dependent anion protein

Key words: apoptosis, human neoplastic cells, lamins, perylenequinones, polyomavirus-transformed rat fibroblasts, reactive oxygen species




Figure 1. The photoactivation of CalC to CalC^{ΦE} sets off the rapid and complete destruction of the survival-essential lamin B1 at the nuclear envelope. (A) The chemical structure of CalC fully justifies its lipophilicity and ability to enter cells and to be enriched in the cytoplasmic membranes (ER and Golgi apparatus) and nuclear envelope. However, photoactivation is necessary for the biological effects of CalC^{ΦE}, like destruction of lamin B1, inhibition of PKCs (IC50 = 50 nM), and apoptosis, to be made happen. If never exposed to light since the onset of the experiments, cell-permeating CalC is totally harmless (14). (B) Lamins A/C and emerlin were not changed by a 30-min exposure to CalC^{ΦE} followed by a 30-90-min incubation in the dark still in the presence of the rapidly inactivated CalC; by neat contrast, lamin B1 was totally destroyed within 60 min by the same treatment; strikingly, lamin B1 was not altered by a 60-120-min direct exposure to CalC but no light (14). (C) On the left, the time-related progressive disappearance of 69-kDa lamin B1 holoprotein and the transient appearance of 210- and 244-kDa predestruction bands recognized by the antilamin B1 antibody in the nuclear membrane fractions (NMFs) of cells exposed for 30 min to CalC^{ΦE} and next incubated in the dark with no medium change; the 0-min band is from illuminated only control cells. Conversely, NMF-associated 69-kDa lamin B1 did not wane and no anti-lamin B1 antibody-recognized 210- and 244-kDa bands obtained in the NMFs when cells were exposed to CalC but no light. In either instance, equal sample loading was monitored by reprobing the immunoblots with the anti-emerlin antibody. The immunofluorescent pictures (B) and the immunoblots (c) are typical of five distinct experiments (14).



first, if not its first, critical target of CalC^{qE}, is the lamina (NL). This is the fibrous network that lines the inner surface of the nuclear envelope (15-20). The lamina consists basically of 3-D bundles of lamins A/C piled on top of a sheet of parallel filaments of isoprenylated lamin B filaments joined at 15-nm intervals along their length by short (e.g., 5-nm), non-continuous cross-linkers (15-20). It anchors the nuclear pores plus various other structural and functional components to the nuclear envelope and its periphery (15-20). It controls the transcription of some genes by sequestering transcription factors such as c-Fos and β -catenin and preventing them from getting to their target genes (21). Isoprenylated lamin B1 is permanently attached to the inner nuclear membrane orienting and stabilizing the nucleus in the cell by attaching it to the surrounding cytoskeleton and, for example, preventing the nucleus from spinning (22). Lamin B is the ancestral lamin that is expressed by all cells at all stages of development and, unlike the more mobile lamins A/C, it is essential for the structural integrity of the nuclear envelope and normal nuclear functioning. Therefore, knocking down lamin B1 and B2 proteins with RNAi apoptotically kills cells such as human cervical carcinoma (HCC) HeLa S6 cells and F5 and FR (wt648) rat fibroblasts, while knocking down lamins A/C displaces the associated emerin, but does not kill the cells (23).

We have recently reported that loading polyomavirus-transformed pyF111 rat fibroblasts as well as HCC C4-I cells, in an *in vitro* model for late-stage invasive human cervical cancer (24), with CalC and producing CalC^{qE} by illuminating them with visible light for 30 min, rapidly and selectively destroys lamin B1 (14). Despite the layering of bundles of lamin A/C on the targeted lamin B1 (17), they are not destroyed (Fig. 1B and C), nor are the lamin A/C-associated emerin and the nucleoplasmic cyclin E affected (14). CalC^{qE} is also apoptogenic (14,25). Surprisingly, apoptogenesis does not start in neoplastic rat fibroblasts and HCC C4-I cells until lamin B1 has been mostly destroyed and CalC^{qE} turned off by transferring the lamin B1-depleted cells in a dark incubator (14).

Here, we review the emerging case for a new multimodal model for CalC photodynamic action consisting not just of PKC inhibition and ER stress, but of light-induced selective lamin B1 destruction and PKC inhibition followed, after transfer to the dark, by rebounding cytoplasmic PKC- β_1 and PKC- δ activities and the onset of apoptogenesis.

2. The light-induced events and their in-dark sequelae

CalC^{qE} rapidly enters the cytoplasm of cells and, like hypericin (6), accumulates mainly in the Golgi apparatus and the ER membranes triggering ER stress (4,5). It does not enter the nucleus, although because of its high lipophilicity it probably also collects in the ultimately ER-derived cell nuclear envelope (18). During the first 30 min of illumination, the nuclear envelope lamin B1 of the CalC^{qE}-loaded transformed rat fibroblasts or HCC C4-I cells undergoes a brisk, progressive decline and totally disappears during the first 30 min of incubation in the dark (where CalC^{qE} is rapidly inactivated to CalC) (14) (Fig. 1B and C). While the lamin B1 is being destroyed, neither nuclear membrane lamins

A/C nor the protein emerin, the lamin A-associated binder of transcription factors and part of the four-protein nuclear motor (containing nuclear actin, emerin, lamin A, and nuclear myosin) that moves chromosomes to reposition genes in response to various signals (21,26), are affected (14). The nucleoplasmic cyclin E is also untouched even though within the first 10 min of illumination the aqueous peroxides resulting from the cytoplasmic-located, short-range firing, ephemeral reactive oxygen species (ROS, $^1\text{O}_2$ and O_2^-) produced by the photoactivated CalC^{qE} significantly increase their levels in the nucleus (14).

During the 30 min of illumination and CalC^{qE} activity in the cytoplasm, cytoplasmic PKC- β_1 and PKC- δ activities rapidly drop close to zero, as expected (Fig. 2A and B) (25,27). There are no indications of incipient apoptogenesis such as mitochondrial cytochrome *c* release or caspase-3 activation (Fig. 2C and D) (14,27). However, after the cells are transferred to the dark, while lamin B1 destruction continues to completion, the photo-suppressed cytoplasmic particulate fraction (CPF)-associated PKC- β_1 and PKC- δ activities rebound within minutes (Fig. 2A and B) (25,27).

The prompt in-dark resumption of PKC activities in the CPF is accompanied by the first sign of apoptogenesis (28), the rapid-onset release of mitochondrial cytochrome *c* (Fig. 2C). This release then slows down between 60 and 120 min and finally levels off between 120 and 240 min (Fig. 2C). The cytochrome *c* release is followed by a steadily increasing activity of executioner caspase-3 starting around 90 min (Fig. 2D) (25), and more than 90% of the cells are killed by 3.5 h after shutting off the light (14,27).

Finally, it must be noted that none of these ultimately lethal events, the destruction of lamin B1 or apoptogenesis, are triggered by incubating the pyF111 cells with 75 nM unexcited CalC in the dark. With CalC, but without CalC^{qE}, the cells keep their normal lamin B1-supported nuclear envelopes, continue proliferating normally and double their numbers by 24 h in the dark (14,27).

3. The drivers of CalC^{qE} cytotoxic actions

The early intranuclear accumulation of aqueous peroxides derived from $^1\text{O}_2$ and O_2^- produced by cytoplasm-located, lipophilic CalC^{qE} might be the cause of the so far inexplicably selective oxidation of the nuclear envelope of lamin B1, whereas the closely placed lamin A or lamin A-associated protein emerin or the intranuclear cyclin E are simultaneously spared (14). The oxidized lamin B1, like any other oxidized proteins (29), is in turn proteolyzed by a stimulated proteasomal peptidyl-glutaminase-like protease (14). The turned-on proteasome has an ongoing inertia. So, turning off the light and with it CalC^{qE} at 30 min does not stop oxidized lamin B1 proteolysis from going to completion in as many minutes (Fig. 1C) (14).

Different PKCs have also been shown to be involved in apoptogenesis as well as proliferation and other activities (25,27,30,31). Therefore, as expected, when CalC^{qE} is driving oxidized lamin B1 destruction, it silences PKC- β_1 and PKC- δ and any indication of impending apoptogenesis (Fig. 2A-D). Thus, for example, as the activities of the PKC- δ and PKC- β_1




Figure 2. The very fast rebounding of the activities of the CPF-linked PKC- β_1 and PKC- δ from the CalC^{qE}-induced transient inhibition after transfer to the dark concurs with the brisk onset of apoptogenic events. (A and B) Exposure to CalC^{qE} elicited within 30 min the virtually total inhibition of immunoprecipitable PKC- β_1 - and PKC- δ -specific activities linked to the CPF. However, soon after CalC^{qE} was inactivated to CalC by transfer to a dark incubator, the CPF-linked immunoprecipitable PKC- β_1 - and PKC- δ -specific activities promptly rebounded. SEMs, not shown, were within $\pm 12\%$ the corresponding mean values (25,27 and previously unpublished). (C) A massive release of cytochrome *c* from mitochondria into the cytosol occurred soon after CalC^{qE} was inactivated to CalC by transfer to a dark incubator, and was concomitant with the rebounding CPF-linked PKC- β_1 - and PKC- δ activities, but preceded the activation of executioner caspase-3. Every point on the curves is the mean value \pm SEM of three distinct experiments, each carried out in triplicate (25). (D) The activation of apoptotic caspase-3 in CalC^{qE}-exposed pyF111 cells occurred well downstream of the total waning of NMF-associated lamin B1. The levels of caspase-3 activity in both the nuclear and cytoplasmic fraction remained unchanged for up to 90 min (30-min exposure to CalC^{qE} + 60-min incubation with dark-inactivated CalC) prior to surging during the following 180 min of staying in the dark. Points on the curves are means from 3-5 independent experiments. SEMs (data not shown) were within $\pm 11\%$ of the mean values (14).

holoenzymes bound to the CPF drop virtually to zero by 30 min. During the first 30 min after CalC^{qE} is turned off in the dark, the PKC- β_1 and PKC- δ activities in the CPF, to which they translocate, surge very sharply upwards (Fig. 2A and B) (27). This sudden and large rebound from PKC- β_1 and PKC- δ inactivity is unexpected because of the alleged irreversibility of PKC inhibition in C6 glioma cells by CalC^{qE} (7). Gopalakrishna *et al* (8) reported that washing CalC out

of the glioma cells did not immediately restore PKC activity. More precisely, re-establishment of PKC activity took 12 h instead of the few minutes it took in the transformed rat fibroblasts (24,27) and thus required the reloading of the glioma cells with newly made PKCs. This might mean either that the photoinactivation of PKCs by CalC^{qE} in the pyF111 cells is in fact reversible or that these cells have a store of inactive non-membrane-associated PKCs that are



SPANDIDOS PUBLICATIONS
cessible to lipophilic CalC^{PE} and its ROS photo-
but can be recruited to cytoplasmic membranes
and there swiftly activated when CalC^{PE} disappears.

4. Conclusions

It is generally believed that CalC^{PE} owes its ability to act as a potential photodynamic chemotherapeutic agent by selectively and irreversibly preventing growth factor- or tumor promoter-stimulated PKCs from driving the proliferation of neoplastic cells (32,33). Since it is highly lipophilic, CalC probably collects in cell membranes where the photoexcited CalC^{PE} can fire ¹O₂ and O₂⁻ onto PKCs that have been recruited there. The bombardment of a membrane-associated PKC with ¹O₂ and O₂⁻ from CalC^{PE} prevents diacylglycerol (DAG) released from the membrane's phospholipids by a receptor-activated phospholipase C or a tumor-promoting phorbol ester from binding to the irreversibly altered DAG/phorbol ester kinase binding site in its C1 domain, which normally would activate the enzyme by causing its pseudosubstrate domain to lift away from, and stop blocking, the catalytic domain (8,33-36).

But such a PKC-blocking-only mechanism appears be only half or even less of the story for at least the neoplastic rat fibroblasts and HCC C4-I cells. Instead, CalC^{PE} is a multiple attacking cytotoxic agent with at least two of its targets being equally lethal. When photoexcited in the cytoplasm, the CalC^{PE}-generated ¹O₂ and O₂⁻ and their hydroperoxide derivatives selectively oxidize and thereby target the nuclear membrane indispensable lamin B1 for seemingly total destruction (14). As expected, at the same time CalC^{PE} inhibits cytoplasmic PKC-β₁ and PKC-δ. But contrary to expectations (8,35), turning off the light promptly unleashes PKC-β₁ and PKC-δ activities in the CPF, which coincides with the release of mitochondrial cytochrome *c* (27) that starts the apoptogenic caspase cascade (28).

A possible CalC^{PE}-inhibitable mechanism by which the surging PKCs might trigger cytochrome *c* release and apoptosis is suggested by Chan *et al* (36). They have shown that treating rat rostral ventrolateral medulla cells with *Escherichia coli* lipopolysaccharide causes the translocation of cytosolic PKC (including PKC-β₁ and PKC-δ)/Bax complexes to cell membranes, where Bax is released and next translocated to mitochondria. There, Bax forms complexes with ANT (adenine nucleotide translocase) or VDAC (voltage-dependent anion protein) that cause the opening of the mitochondrial permeability transition pore, the release of the apoptogenic cytochrome *c* into the cytoplasm, and the activation of the apoptogenic caspases.

Clearly other perylenequinones should now be studied to see if this remarkable multimodal lethal capability of CalC^{PE} for neoplastic cells is shared by all the members of this family of photodynamic therapeutic agents.

Acknowledgements

This work was supported in part by the Italian MIUR (Ministero dell'Istruzione, dell'Università e della Ricerca,

ex-60% allotment funds) and by MIUR-PRIN (Cofin) funds.

References

1. Beck TP, Kirsh EJ, Chmura SJ, Kovar DA, Chung T, Rinker-Schaeffer CW and Stadler WM: *In vitro* evaluation of calphostin C as a novel agent for photodynamic therapy of bladder cancer. *Urology* 54: 573-577, 1999.
2. Diwu Z and Lown JW: Photosensitization with anticancer agents. 15. Perylenequinoid pigments as potential photodynamic therapeutic agents: formation of semiquinone radicals and reactive oxygen species on illumination. *J Photochem Photobiol B* 18: 131-143, 1993.
3. Gomer CJ: Preclinical examination of first and second generation photosensitizers used in photodynamic therapy. *Photochem Photobiol* 54: 1093-1107, 1991.
4. Kaul A and Maltese WA: Killing of cancer cells by the photo-activatable protein kinase C inhibitor, calphostin C, involves induction of endoplasmic reticulum stress. *Neoplasia* 11: 823-834, 2009.
5. Olivo M and Ali-Sayed M: Apoptosis signaling mechanisms in human cancer cells induced by calphostin-PDT. *Int J Oncol* 30: 537-548, 2007.
6. Sanovic R, Krammer B, Grumböck S and Verwanger T: Time-resolved gene expression profiling of human squamous cell carcinoma cells during the apoptosis process induced by photodynamic treatment with hypericin. *Int J Oncol* 35: 921-939, 2009.
7. Bruns RF, Miller FD, Merriman RL, Howbert JJ, Heath WF, Kobayashi E, Takahashi I, Tamaoki T and Nakano H: Inhibition of protein kinase C by calphostin C is light-dependent. *Biochem Biophys Res Commun* 176: 288-293, 1991.
8. Gopalakrishna R, Chen ZH and Gundimeda U: Irreversible oxidative inactivation of protein kinase C by photosensitive inhibitor calphostin C. *FEBS Lett* 314: 149-154, 1992.
9. Kobayashi E, Ando K, Nakano H and Tamaoki T: UCN-1028A, a novel and specific inhibitor of protein kinase C from *Cladosporium*. *J Antibiotics (Tokyo)* 42: 153-155, 1989.
10. Kobayashi E, Nakano H, Morimoto M and Tamaoki T: Calphostin C (UCN-1028c), a novel microbial compound, is a highly potent and specific inhibitor of protein kinase C. *Biochem Biophys Res Commun* 159: 548-553, 1989.
11. Pollack IF and Kawecky S: The effect of calphostin C, a potent photodependent protein kinase C inhibitor, on the proliferation of glioma cells *in vitro*. *J Neurooncol* 31: 255-266, 1997.
12. Takao I, Kobayashi E, Yoshida M and Sano H: Calphostins, novel and specific inhibitors of protein kinase C. B. Chemical structures. *J Antibiotics (Tokyo)* 42: 1475-1481, 1989.
13. Gueded RC and Eriksson LA: Photophysics, photochemistry, and reactivity: molecular aspects of perylenequinone reactions. *Photochem Photobiol Sci* 6: 1089-1096, 2007.
14. Chiarini A, Whitfield JF, Pacchiana R, Armato U and Dal Pra I: Photoexcited calphostin C selectively destroys nuclear lamin B1 in neoplastic human and rat cells. A novel mechanism of action of a photodynamic tumor therapy agent. *Biochim Biophys Acta* 1783: 1642-1653, 2008.
15. Bridger JM, Foeger N, Kill LR and Hermann H: The nuclear lamina. Both a structural framework and a platform for genome organization. *FEBS J* 274: 1354-1361, 2007.
16. Dechat T, Pflieger K, Sengupta K, Shimi T, Schumaker DK, Solimando L and Goldman RD: Nuclear lamins: major factors in the structural organization and function of the nucleus and chromatin. *Genes Dev* 22: 832-853, 2008.
17. Goldberg MW, Fiserova J, Huttenlauch I and Stick R: A new model for nuclear lamina organization. *Biochem Soc Trans* 36: 1339-1343, 2008.
18. Larijani B and Poccia DL: Nuclear envelope formation: mind the gaps. *Annu Rev Biophys* 38: 107-124, 2009.
19. Goldman RD, Gruenbaum Y, Moir RD, Schumaker DK and Spann TP: Nuclear lamins: building blocks of nuclear architecture. *Genes Dev* 16: 533-547, 2002.
20. Gruenbaum Y, Margalit A, Goldman RD, Schumaker DK and Wilson KL: The nuclear lamin comes of age. *Nat Rev Mol Cell Biol* 6: 21-31, 2005.
21. Heessen S and Fornerod M: The inner nuclear envelope as a transcription factor resting place. *EMBO Rep* 8: 914-919, 2007.

22. Ji JY, Lee RT, Vergnes L, Fong LG, Stewart CL, Reue K, Young SG, Zhang Q, Shanahan CM and Lammerding J: Cell nuclei spin in the absence of lamin B1. *J Biol Chem* 282: 20015-20026, 2007.
23. Harborth J, Elbashir SM, Bechert K, Tuschl T and Weber K: Identification of essential genes in cultured mammalian cells using small interfering RNAs. *J Cell Sci* 114: 4557-4565, 2001.
24. Carlson MW, Iyer VR and Marcotte EM: Quantitative gene expression assessment identifies appropriate cell line models for individual cervical cancer pathways. *BMC Genomics* 8: 117, 2007.
25. Dal Pra I, Whitfield JF, Chiarini A and Armato U: Increased activity of the protein kinase- δ holoenzyme in the cytoplasmic particulate fraction precedes activation of caspases in the polyomavirus-transformed pyF111 rat fibroblasts exposed to calphostin C or topoisomerase II inhibitors. *Exp Cell Res* 255: 171-183, 2000.
26. Mehta IS, Elcock LS, Amira M, Kill IR and Bridger JM: Nuclear motors and nuclear structures containing A-type lamins and emerin: is there a functional link? *Biochem Soc Trans* 36: 1384-1388, 2008.
27. Chiarini A, Whitfield JF, Armato U and Dal Pra I: VP-16 (etoposide) and calphostin C trigger different nuclear but akin cytoplasmic patterns of changes in the distribution and activity of protein kinase C- β_1 in polyomavirus-transformed pyF111 rat fibroblasts. *Int J Mol Med* 17: 111-120, 2006.
28. Yan N and Shi Y: The mechanisms of apoptosis through structural biology. *Annu Rev Cell Dev Biol* 21: 35-56, 2005.
29. Jung T and Grune T: The proteasome and its role in the degradation of oxidized proteins. *IUBMB Life* 60: 743-752, 2008.
30. Chiarini A, Whitfield JF, Armato U and Dal Pra I: Protein kinase C- β_{II} is an apoptotic lamin kinase in polyomavirus-transformed etoposide-treated pyF111 rat fibroblasts. *J Biol Chem* 277: 18827-18839, 2002.
31. Reyland ME: Protein kinase C- δ and apoptosis. *Biochem Soc Trans* 35 (Pt. 5): 1001-1004, 2007.
32. Da Rocha AB, Mans DRA, Regner A and Schwartzmann G: Targeting protein kinase C: new therapeutic opportunities against high-grade malignant gliomas? *Oncologist* 7: 17-33, 2002.
33. Whitfield JF and Chakravarthy B: Calcium: The Grand-Master Cell Signaller. NRC Research Press, Ottawa, 2001.
34. Newton AC: Protein kinase C: structure, function and regulation. *J Biol Chem* 270: 28495-28498, 1995.
35. Rotenberg SA, Huang MH, Zhu J, Su L and Riedel H: Deletion analysis of protein kinase C inactivation by calphostin C. *Mol Carcinogenesis* 12: 42-49, 1995.
36. Chan JY, Chang AY, Wang LL, Ou CC and Chan SH: Protein kinase C-dependent mitochondrial translocation of pro-apoptotic protein Bax on activation of inducible nitric-oxide synthase in rostral ventrolateral medulla mediates cardiovascular depression during experimental endotoxemia. *Mol Pharmacol* 71: 1129-1139, 2007.