

Table SII. Adverse event characteristics in patients with regorafenib-induced hepatic toxicity.

| Patient no. | Baseline parameters CTCAE grade | Maximum CTCAE grade | Pattern |
|-------------|--|--|----------------|
| 1 | ALT: 0, AST: 0, AP: 2, γ -GT: 3, Bili: 1 | ALT: 3, AST: 4, AP: 2, γ -GT: 3, Bili: 2 | Hepatocellular |
| 2 | ALT: 0, AST: 0, AP: 0, γ -GT: 0, Bili: 0 | ALT: 0, AST: 0, AP: n.a., γ -GT: n.a., Bili: 1 | Cholestatic |
| 3 | ALT: 1, AST: 1, AP: 2, γ -GT: 3, Bili: 2 | ALT: 1, AST: 1, AP: 2, γ -GT: 3, Bili: 3 | Mixed |
| 4 | ALT: 0, AST: 1, AP: 0, γ -GT: 0, Bili: 0 | ALT: 4, AST: 4, AP: 1, γ -GT: 3, Bili: 3 | Mixed |
| 5 | ALT: 0, AST: 0, AP: 0, γ -GT: 0, Bili: 0 | ALT: 2, AST: 2, AP: 1, γ -GT: 3, Bili: 3 | Mixed |

CTCAE, Common Terminology Criteria of Adverse Events; ALT, alanine aminotransferase; AST, aspartate aminotransferase; AP, alkaline phosphatase; γ -GT, γ -glutamyltransferase; Bili, bilirubin; n.a., not available.