

Table SI. Primers used in the study for qPCR (NCBI-blast).

Gene	Forward primer (5'-3')	Reverse primer (5'-3')	Length (bp)
<i>LRPPRC</i>	GAGAGATGCCGGAATTGAGC	CTCGGACTTCTCCACCTTCT	118
<i>ABCB1</i>	CACCCGACTTACAGATGATG	GTTGCCATTGACTGAAAGAA	81
<i>MRP1</i>	GCCGAAGGAGAGATCATC	AACCCGAAAACAAAACAGG	109
<i>BCRP</i>	TGACGGTGAGAGAAAACCTTAC	TGCCACTTTATCCAGACCT	122
<i>LRP</i>	GGCAGGACAATGAGAGGGTACT	CCGAACTTGCCCTGTGACAT	137
<i>GAPDH</i>	AATCCCATCACCATCTTCCA	TGGACTCCACGACGTACTCA	82

LRPPRC, leucine-rich PPR-motif-containing protein; *ABCB1*, ATP Binding Cassette Subfamily B Member 1; *MDR1*, multidrug resistance-associated protein 1; *BCRP*, breast cancer-resistant protein; *LRP*, lung resistance-related protein; *GAPDH*, glyceraldehyde 3-phosphate dehydrogenase.