

Table SI. The scores of each soft tissue tumor with 3 or more lesions, according to the original scoring system.

|                                      | n  | Mean $\pm$ SD | 95% CI   | Min | Max |
|--------------------------------------|----|---------------|----------|-----|-----|
| Benign                               |    |               |          |     |     |
| Lipoma                               | 35 | 0.6 $\pm$ 0.6 | 0.4-0.8  | 0   | 2.5 |
| Schwanoma                            | 23 | 2.1 $\pm$ 1.3 | 1.5-2.6  | 0   | 4.5 |
| Vascular malformation                | 9  | 1.2 $\pm$ 1.3 | 0.3-2.2  | 0   | 4   |
| Epidermal cyst                       | 9  | 0.2 $\pm$ 0.4 | -0.1-0.6 | 0   | 1   |
| Spindle cell lipoma                  | 5  | 1.9 $\pm$ 1.8 | -0.4-4.2 | 0   | 4   |
| Tenosynovial giant cell tumor        | 3  | 0.0 $\pm$ 0.0 | 0.0-0.0  | 0   | 0   |
| Intermediate                         |    |               |          |     |     |
| Atypical lipomatous tumor            | 8  | 1.4 $\pm$ 1.1 | 0.5-2.3  | 1   | 4   |
| Desmoid type fibromatosis            | 5  | 3.8 $\pm$ 1.6 | 1.9-5.7  | 1.5 | 5.5 |
| Plantar fibromatosis                 | 4  | 0.8 $\pm$ 0.9 | -0.6-2.1 | 0   | 1.5 |
| Malignancy                           |    |               |          |     |     |
| Myxofibrosarcoma                     | 14 | 3.8 $\pm$ 1.5 | 2.9-4.6  | 1   | 5.5 |
| Myxoid liposarcoma                   | 5  | 4.0 $\pm$ 1.5 | 2.1-5.9  | 2.5 | 5.5 |
| Undifferentiated pleomorphic sarcoma | 4  | 5.0 $\pm$ 0.6 | 4.1-5.9  | 4.5 | 5.5 |
| Soft tissue metastasis of carcinoma  | 4  | 3.8 $\pm$ 1.5 | 1.4-6.1  | 1.5 | 4.5 |
| Rhabdomyosarcoma                     | 3  | 4.7 $\pm$ 0.8 | 2.8-6.6  | 4   | 5.5 |
| Leiomyosarcoma                       | 3  | 4.8 $\pm$ 0.6 | 3.4-6.3  | 4.5 | 5.5 |