

Table SI. Final PD score of the peritoneal dissemination model for immune cell analysis (on Days 6, 13, 20).

Cell line used	PD Score ^a			
	P0	P1	P2	P3
MC38 (n=18)				
Day 6	4	2	0	0
Day 13	1	1	2	2
Day 20	0	0	3	3
CT26 (n=12)				
Day 6	0	2	1	1
Day 13	0	0	1	3
Day 20	0	0	0	4

^aPD score: P0 for no nodules, P1 for 1-3 nodules, P2 for 4-9 nodules, and P3 for 10 or more nodules. PD, peritoneal dissemination.

Table SII. Final tumor volume of the subcutaneous inoculated mice for immune cell analysis (on Days 6, 13, 20) and survival analysis.

Day	Maximum diameter (mm)	Minimum diameter (mm)	V (mm ³) ^b
Day 6 (n=6) ^a	4.4±1.4	3.6±1.0	34.3±26.2
Day 13 (n=6) ^a	7.1±1.5	5.5±0.8	114.6±47.8
Day 20 (n=6) ^a	11.1±1.9	9.3±1.4	508.6±216.8
Survival analysis (n=6) ^a	22.2±1.5	19.1±1.4	4,084.4±845.2

^aData are expressed as the average ± SD. ^bTumor volume (V)=maximum diameter x (minimum diameter)²/2.