

Table SI. Demographic and clinical parameters of patients with DKD.

Variables	Control group (n=10)	Micro group (n=10)	Macro group (n=10)	P-value
Age, years	32.4±6.3	65.2±16.2	59.1±11.58	P<0.01 <sup>a,b</sup>
SBP, mmHg	118.8±9.6	140±21.9	142.3±19.0	P<0.01 <sup>a,b</sup>
BUN, mmol/l	4.58±0.94	7.04±2.67	11.47±4.18	P<0.01 <sup>a</sup>
Scr, μmol/l	74.9±12.1	90.8±28.9	278.9±214.9	P<0.01 <sup>a</sup>
FPG, mmol/l	4.78±0.45	7.67±4.12	8.44±3.36	P<0.01 <sup>a,b</sup>
HbA1c, %	5.09±0.65	9.92±2.3	7.82±2.58	P<0.01 <sup>a,b</sup>
LDL-C, mmol/l	2.60±0.45	2.66±1.12	3.20±1.85	NS
HDL-C, mmol/l	1.50±0.35	1.11±0.30	1.16±0.43	P<0.01 <sup>a,b</sup>

<sup>a</sup>Macro group vs. Control group; <sup>b</sup>Micro group vs. Control group. Micro, DKD-microalbuminuria (UACR, 30-300 mg/g); Macro, DKD-macroalbuminuria (UACR, >300 mg/g); SBP, systolic blood pressure; BUN, blood urea nitrogen ; Scr, serum creatinine; FPG, fasting plasma glucose; HbA1c, hemoglobin A1C; LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol; NS, not significant. Data are presented as the mean ± standard deviation.