

Table S1. Primer sequences used for reverse transcription-quantitative PCR.

Gene	Forward (5'-3')	Reverse (5'-3')
HDAC 1	ACCGGGCAACGTTACGAAT	CTATCAAAGGACACGCCAAGTG
HDAC 2	TCATTGGAAAATTGACAGCATAGT	CATGGTGATGGTGTGAAGAAG
HDAC 3	TTGAGTTCTGCTCGCGTTACA	CCCAGTTAATGGCAATATCACAGAT
HDAC 4	AATCTGAACCACTGCATTTCCA	GGTGGTTATAGGAGGTGCGACT
HDAC 5	TTGGAGACGTGGAGTACCTTACAG	GACTAGGACCACATCAGGTGAGAAC
HDAC 6	TGGCTATTGCATGTTCAACCA	GTCGAAGGTGAACTGTGTTCTT
HDAC 7	CTGCATTGGAGGAATGAAGCT	CTGGCACAGCGGATGTTTG
HDAC 8	TCCCGAGTATGTCAGTATATATGA	GCTTCAATCAAAGAATGCACCA
HDAC 9	CAGGCGGAAGGATGGAAATG	ATGCGTTGCTGTGAAACCAT
HDAC 10	GGCCTTTGAGTTTGACCCTG	CAGCGTCTGTACTGTCATGC
HDAC 11	TGTCTACAACCGCCACATCT	CGGTGCCTGCATTGTATACC
GAPDH	GCTCATTTCTGGTATGAC	ACAGGGTACTTTATTGATGGT
Ki-67	CGGATCGTCCCAGTGGAAG	TTGCCTCCTGCTCATGGATT
E2Ftranscriptionfactor	GCTGGAGCTAGGAGAAAGCG	CAAGAGACGTATCATACCGCGT
18SrRNA	GATCAAACCAACCCGGTCA	CCGTTTCTCAGGCTCCCTCT

HDAC, histone deacetylase; E2F3,