

Table S1. Histological score.

Feature graded	Description	Grade
Inflammation	None	0
	Slight	1
	Moderate	2
	Severe	3
Extent	None	0
	Mucosa	1
	Submucosa	2
	Transmural	3
Percentage of involvement, %	1-25	0
	26-50	1
	51-75	2
	76-100	3
Crypt damage	None	0
	Submucosa	1
	Basal one-thirds lost	2
	Basal two-thirds lost	3
	Only surface epithelium intact	4
	All crypts and epithelium are destroyed	5
Ulceration	No ulceration	0
	Mild ulceration	1
	Moderate ulceration	2
	Extensive ulceration	3

Table SII. PCR primer sequences.

Genes	Primers
GPR32	Forward: 5'-TGTGGTTATCCTGTCTGCGTC-3'
	Reverse: 5' -GGCCAATGGTCCCTATAATGT-3'
FPR2	Forward: 5'-GCCTTTTGGCTGGTTCCTGTGT-3'
	Reverse: 5' -GCCTTTTGGCTGGTTCCTGTGT-3'
ChemR23	Forward: 5'-CTCACTCCCAAATGGACCCTGT-3'
	Reverse: 5' -CACACGATGGTGAGGTAGCAAG-3'
TNF- α	Forward: 5'-CTCTTCTGCCTGCTGCACTTTG-3'
	Reverse: 5' -ATGGGCTACAGGCTTGTCACCTC-3'
IL-1 β	Forward: 5'-TGCCACCTTTTGACAGTGATG-3'
	Reverse: 5' -TGATACTGCCTGCCTGAAGC-3'
IL-6	Forward: 5'-AGACAGCCACTCACCTCTTCAG-3'
	Reverse: 5' -TTCTGCCAGTGCCTCTTTGCTG-3'
COX2	Forward: 5'-GAAGTCTTTGGTCTGGTGCCTG-3'
	Reverse: 5' -GTCTGCTGGTTTGGAATAGTTGC-3'
iNOS	Forward: 5'-GTTCTCAGCCCAACAATACAAGA-3'
	Reverse: 5' -GTGGACGGGTCGATGTCA-3'
β -actin	Forward: 5'-CTACAGCTTCACCACCACAG-3'
	Reverse: 5'-CTACAGCTTCACCACCACAG-3'

GPR32, G protein-coupled receptor 32; FPR2, formyl peptide receptor 2; ChemR23, chemokine like receptor 1; iNOS, inducible nitric oxide synthase; COX2, cytochrome c oxidase subunit 2.