

Table SI. Relationship between cancer stage and NE or ALB.

Factor	Total, n=77	Stage 0, n=11	Stage I, n=14	Stage II, n=22	Stage III, n=17	Stage IV, n=13	P-value
NE, median (IQR)	3.37 (2.68, 4.83)	4.19 (3.04, 5.72)	3.50 (3.14, 4.86)	3.24 (2.74, 3.82)	3.15 (2.46, 5.04)	3.77 (2.73, 4.96)	0.0518
ALB, mean ± SD	43.37±4.50	45.07±3.78	43.75±5.28	43.83±4.16	42.12±5.41	42.36±3.21	0.0431

NE, neutrophils; ALB, albumin.

Table SII. Relationship between cancer stage and NE or ALB before and after chemotherapy.

Factor	Total, n=77	Stage II, n=22	Stage III, n=17	P-value
Before chemotherapy				
NE, Median (IQR)	3.37 (2.68, 4.83)	3.24 (2.74, 3.82)	3.15 (2.46, 5.04)	0.531
ALB, Mean \pm SD	43.37 ± 4.50	43.83 ± 4.16	42.12 ± 5.41	0.621
After chemotherapy				
NE, Median (IQR)	3.46 (2.49, 4.60)	3.05 (2.46, 4.08)	3.44 (2.10, 4.94)	0.683
ALB, Median (IQR)	42.63 (39.60, 44.90)	42.72 (41.94, 43.80)	39.90 (38.81, 40.67)	<0.001 ^a

^aP<0.001. NE, neutrophils; ALB, albumin.