

Table SI. Overall survival univariate Cox regression analysis.

Variables	B	SE	df	P-value	Exp(B)	95% CI for Exp(B)	
						Lower	Upper
Age (≤ 50 vs. > 50 years)	0.218	0.543	1	0.688	1.243	0.429	3.602
Sex (male vs. female)	-0.523	0.393	1	0.183	0.593	0.275	1.280
Performance status ^a			2	0.035			
1 vs. 0	9.171	127.175	1	0.943	9614.878	0.000	1.717×10^{112}
2 vs. 0	10.373	127.176	1	0.935	31996.263	0.000	5.719×10^{112}
Tumour stage ^b			3	0.810			
2 vs. 1	-0.029	0.450	1	0.948	0.971	0.402	2.345
3 vs. 1	0.233	0.774	1	0.763	1.263	0.277	5.762
4 vs. 1	0.471	0.544	1	0.387	1.602	0.551	4.654
Nodal stage ^b			2	0.222			
2 vs. 1	0.463	0.435	1	0.288	1.589	0.677	3.730
3 vs. 1	0.872	0.528	1	0.099	2.392	0.849	6.736
Brachytherapy (yes vs.no)	-0.270	0.614	1	0.661	0.764	0.229	2.544
Composite stage ^b			3	0.333			
II vs. I	-0.662	0.667	1	0.321	0.516	0.140	1.907
III vs. I	0.254	0.505	1	0.616	1.289	0.479	3.468
IV vs. I	0.537	0.490	1	0.273	1.712	0.655	4.470
Type of treatment			2	0.831			
EBRT vs. brachytherapy	0.334	0.631	1	0.597	1.396	0.405	4.808
Surgery vs. brachytherapy	0.153	0.678	1	0.821	1.166	0.308	4.405

^aUsing the Eastern Cooperative Oncology Group scale (28); ^busing the American Joint Committee on Cancer Staging Manual (12). CI, confidence interval; df, degrees of freedom; EBRT, external beam radiotherapy; SE, standard error.

Table SII. Disease-free survival univariate Cox regression analysis.

Variables	B	SE	df	P-value	Exp(B)	95% CI for Exp(B)	
						Lower	Upper
Age (≤ 50 vs. > 50 years)	-0.300	0.357	1	0.401	0.741	0.368	1.491
Sex (male vs. female)	-0.365	0.289	1	0.207	0.694	0.394	1.223
Performance status ^a			2	0.381			
1 vs. 0	0.053	1.013	1	0.958	1.055	0.145	7.688
2 vs. 0	0.590	1.062	1	0.579	1.804	0.225	14.472
Tumour stage ^b			3	0.025 ^c			
2 vs. 1	0.630	0.357	1	0.078	1.877	0.932	3.778
3 vs. 1	0.380	0.643	1	0.554	1.463	0.415	5.156
4 vs. 1	1.201	0.395	1	0.002 ^c	3.323	1.531	7.212
Nodal stage ^b			2	0.005 ^c			
2 vs. 1	0.587	0.322	1	0.068	1.799	0.956	3.384
3 vs. 1	1.263	0.393	1	0.001 ^c	3.538	1.638	7.640
Brachytherapy (yes vs.no)	0.041	0.409	1	0.920	1.042	0.467	2.322
Composite stage ^b			3	0.006 ^c			
II vs. I	0.251	0.450	1	0.577	1.285	0.532	3.105
III vs. I	0.347	0.438	1	0.428	1.415	0.600	3.337
IV vs. I	1.209	0.383	1	0.002 ^c	3.350	1.581	7.099
Type of treatment			2	0.468			
EBRT vs. brachytherapy	0.106	0.422	1	0.802	1.112	0.486	2.542
Surgery vs. brachytherapy	-0.306	0.469	1	0.515	0.737	0.294	1.849

^aUsing the Eastern Cooperative Oncology Group scale (28); ^busing the American Joint Committee on Cancer Staging Manual (12); ^cP<0.05 indicates a statistically significant difference. CI, confidence interval; df, degrees of freedom; EBRT, external beam radiotherapy; SE, standard error.

Table SIII. Disease-free survival multivariate Cox regression analysis.

Variables	B	SE	df	P-value	Exp(B)	95% CI for Exp(B)	
						Lower	Upper
Nodal stage ^a			2	0.005 ^b			
2 vs. 1	0.587	0.322	1	0.068	1.799	0.956	3.384
3 vs. 1	1.263	0.393	1	0.001 ^b	3.538	1.638	7.640

^aUsing the American Joint Committee on Cancer Staging Manual (12); ^bP<0.05 indicates a statistically significant difference. CI, confidence interval; df, degrees of freedom; EBRT, external beam radiotherapy; SE, standard error.