

Table SI. Sequences of primers used for RT-qPCR in the present study.

Gene	Primer sequence (5'-3')
PVT1	GGGGAATAACGCTGGTGGAA
	CCCATGGACATCCAAGCTGT
hsa-miR-152-3p	CGCGTCAGTGCATGACAGA
	AGTGCAGGGTCCGAGGTATT
hsa-miR-23b-3p	CGATCACATTGCCAGGGAT
	AGTGCAGGGTCCGAGGTATT
hsa-miR-27b-3p	GCGCGTTCACAGTGGCTAAG
	AGTGCAGGGTCCGAGGTATT
BLM	GAGTCTGCGTGCGAGGATTA
	AGTGTTCTGGCTGAGTGACG
c-Myc	CTCCGAGATAGCAGGGGACT
	GGGCTTCGCTTACCAGAGT
U6	AAAGCAAATCATCGGACGACC
	GTACAACACATTGTTTCCTCGGA
GAPDH	GGAGCGAGATCCCTCCAAAAT
	GGCTGTTGTCATACTTCTCATGG

PVT1, plasmacytoma variant translocation 1; BLM, bloom syndrome protein.

Table SII. Sequences of shRNAs against specific targets.

shRNA	Sequence (5'-3')
METTL3-sh1	GCTGCACTTCAGACGAATTAT
METTL3-sh2	GCCAAGGAACAATCCATTGTT
PVT1-sh1	GAACTCCTGACCTCAAGTGAT
PVT1-sh2	CGGCACATTCAGGATACTAAAA

PVT1, plasmacytoma variant translocation 1; methyltransferase 3.