

Table SI. The association between clinicopathological characteristics of unresectable GC patients and NLR variations obtained before nivolumab therapy.

Characteristic	NLR		P-value
	Low (N= 14)	High (N=10)	
Sex			0.68
Male	8 (57)	4 (40)	
Female	6 (43)	6 (60)	
Age			0.41
<70 years	5 (35)	6 (60)	
≥70 years	9 (65)	4 (40)	
ECOG performance status			0.23
0	4 (29)	6 (60)	
1	8 (57)	2 (20)	
2	2 (14)	2 (20)	
3	0 (0)	0 (0)	
Primary tumor location			0.03
Upper	3 (21)	7 (70)	
Middle	6 (42)	3 (30)	
Low	5 (37)	0 (0)	
HER2 status			0.62
Positive	3 (21)	1 (10)	
Negative	11 (79)	9 (90)	
Lauren classification			0.68
Intestinal type	6 (43)	6 (60)	
Diffuse type	8 (57)	4 (40)	
Number of metastatic organs			1.00
≤1	7 (50)	5 (50)	
>2	7 (50)	5 (50)	
Liver metastasis			0.67
Yes	4 (29)	4 (40)	
No	10 (71)	6 (60)	
Peritoneal dissemination			0.42
Yes	7 (50)	3 (30)	
No	7 (50)	7 (70)	
Ascites			0.70
Yes	7 (50)	4 (40)	
No	7 (50)	6 (60)	