

Table SI. Patients' characteristics before and after propensity score matching.

Characteristic	Before propensity score matching			After propensity score matching		
	RT ⁻ (n=71)	RT ⁺ (n=41)	SD	RT ⁻ (n=23)	RT ⁺ (n=23)	SD
Age, years	45.0 (22.0, 74.0)	46.0 (28.0, 74.0)	0.039	44.0 (28.0, 71.0)	46.0 (28.0, 68.0)	0.196
FIGO stage IB	63 (88.7)	36 (87.8)	0.029	19 (82.6)	19 (82.6)	<0.001
Histology SCC	38 (53.5)	27 (65.9)	0.253	14 (60.9)	13 (56.5)	0.088
≥4 cm tumor size	15 (21.1)	28 (68.3)	1.078	10 (43.5)	10 (43.5)	<0.001
≥1/3 SI	34 (47.9)	39 (95.1)	1.228	21 (91.3)	21 (91.3)	<0.001
LVSI ⁺	23 (32.4)	25 (61.0)	0.598	14 (60.9)	14 (60.9)	<0.001

Values are expressed as n (%) or median (interquartile range). RT⁺, patients with adjuvant radiotherapy; RT⁻, patients without adjuvant RT; SD, standardized difference; FIGO, International Federation of Obstetrics and Gynecology; SCC, squamous cell carcinoma; SI, stromal invasion; LVSI⁺, patients with lymphovascular space invasion.