

Table SII. Primer sequences used for reverse transcription-quantitative PCR.

Gene	Forward	Reverse
<i>FAS</i>	GGACCCTCCTACCTCTGGTT	ATGATGCAGGCCTTCCAAGT
<i>BAX</i>	GGGGACGAACTGGACAGTAA	GTCCAATGTCCAGCCCATGA
<i>BIM</i>	GTGTGACCGAGAAGGTAGACA	TCTGTCTGTAGGGAGGTAGGG
<i>BCL-2</i>	GAAGTGGGGGAGGATTGTGG	GCCGGTTCAGGTACTIONCAGTC
<i>BCL-XL</i>	CCCAGAAGGGACTGAATCGG	GTGATGTGGAGCTGGGATGT
<i>KIF1B-α</i>	CAGAGACTGGACGCGGATTC	GGTTTGCAACTTGCTGGGAG
<i>KIF1B-β</i>	GAAGAAGTTCCTTGGACACAGC	GGCATTGGCCTCCTTTAGGT
<i>Total KIF1B</i>	GAGGACCCCAATGCCAAACT	GGGACCCCATGGATGTATCA
<i>RBFOX2</i>	GAGCCGACAACAACCTCCTGA	GGCTTGCGTACTTCCGTAGA
<i>GAPDH</i>	GCACCGTCAAGGCTGAGAAC	TGGTGAAGACGCCAGTGGA