

Table S1. Sequences of primers used in reverse transcription-quantitative PCR.

Gene name	Primer sequence, 5'-3'
COL2A1	F: CCCCAGGTTGACAAAG
	R: AGGGATTCCATTAGAGCC
IL-1 β	F: TCTGTGACTCGTGGGA
	R: TGTTGGCTTATGTTCTGT
IL-6	F: AGTTCCGTTTCTACCTG
	R: TCCTTAGCCACTCCTT
DUSP26	F: CTC AACGCCTCCCACAG
	R: CTC ACTCCCACCGCACA
COL1A1	F: CGAGTATGGAAGCGAAGGT
	R: CCACAAGCGTGCTGTAGGT
TNF- α	F: TTCTCATTCTGCTCGTGG
	R: CCTCCGCTTGGTGGTTT
HDAC1	F: GTTCTTGCGTTCTATTCTG
	R: GTCCGTCTGCTGCTTAT
HDAC2	F: GGATGGAGAAGACCCTG
	R: CGACATTCCTACGACCT
HDAC8	F: CCAGAAGGTCAGCCAAGA
	R: TTTCCGTCGCAATCGTA
β -actin	F: TGGCACCACACTTTCTACAATGAGC
	R: GGGTCATCTTTTCACGGTTGG

COL2A1, type II collagen; IL, interleukin; DUSP26, dual-specificity phosphatase 26; COL1A1, type I collagen; HDAC, histone deacetylase; F, forward; R, reverse.