

Table SII. Primers for reverse transcription-quantitative PCR.

Name	Direction	Sequence, 5'→3'
GAPDH	Forward	CCGCATCTTCTTGTGCAGTG
	Reverse	ACCAGCTTCCCATTCTCAGC
CTGF	Forward	ATGGTGCACCCTGTGTCTTC
	Reverse	TGGCTCGCATCATAGTTGGG
Collagen I	Forward	ACATGTTCAGCTTTGTGGACC
	Reverse	CTTGCATAGCACGCCATCG
TGF- $\beta$ 1	Forward	CCCCTACATTTGGAGCCTGG
	Reverse	TTGCGACCCACGTAGTAGAC
CCL20	Forward	TGCTGCCTCACGTACACAAA
	Reverse	CAGCGCACACGGATCTTTTC
CSF1	Forward	AGCAGTTGATCGACAGCCAA
	Reverse	ACACAGGCCTCGTTCTGTTC
CSF2	Forward	AACCCCGGAAACTGACTGTG
	Reverse	CTCCTCATTCTGGACCGGC

CTGF, connective tissue growth factor; TGF- $\beta$ 1, transforming growth factor- $\beta$ 1; CCL20, C-C motif chemokine ligand 20; CSF1, colony-stimulating factor 1.