Adipocyte differentiation of human marrow mesenchymal stem cells reduces the supporting capacity for hematopoietic progenitors but not for severe combined immunodeficiency repopulating cells

NOBUHIKO OOKURA¹, YOSHIHIRO FUJIMORI², KEISUKE NISHIOKA¹, SHUNRO KAI^{2,3},
HIROSHI HARA² and HIROYASU OGAWA^{1,2}

¹Division of Hematology, Department of Internal Medicine, ²Laboratory of Cell Transplantation, Institute for Advanced Medical Sciences, ³Department of Transfusion Medicine, Hyogo College of Medicine, 1-1 Mukogawa-cho, Nishinomiya, Hyogo 663-8501, Japan

Received November 10, 2006; Accepted December 14, 2006

Abstract. Bone marrow stromal cells provide a micro-environment for hematopoiesis. Adipocytes are the major stromal cell phenotype in bone marrow, but their function in hematopoiesis is poorly understood. In this study, we compared the hematopoietic-supporting capacity of adipocytes and their progenitor, mesenchymal stem cells (MSCs), by culturing human cord blood (CB) CD34⁺CD38⁻ hematopoietic progenitor cells (HPCs) on a layer of adipocytes or MSCs. CB CD34⁺CD38⁻ cells cultured on MSCs generated higher proportions of CD34⁺CD38⁻ HPCs and colony-forming cells than those cultures on a layer of adipocytes, indicating an inferior hematopoietic support by adipocytes. However, CB CD34⁺CD38⁻ HPCs cultured on MSCs and adipocytes were equally capable of reconstituting human hematopoiesis in non-obese diabetic/severe combined immunodeficient disease (NOD/SCID) mice. These findings show that differentiation of MSCs into adipocytes is accompanied by the loss of capacity to support mature HPCs, but not transplantable SCID-repopulating cells.

Introduction

Human bone marrow contains hematopoietic stem cells (HSCs) and their progeny (hematopoietic progenitor cells, HPCs) which continuously provide blood cells in the periphery, and stromal cells which provide a supportive scaffolding or microenvironment for hematopoiesis (1).

Human CD34⁺ cells include several classes of HPCs, such as relatively mature *in vitro* colony-forming cells (CFCs) and immature transplantable SCID-repopulating cells (SRC) that can engraft in non-obese diabetic/severe combined immunodeficient disease (NOD/SCID) mice (2,3). Bone marrow stromal cells comprise a variety of cell types, such as macrophages, fibroblasts, endothelial cells, adipocytes, reticular cells and osteogenic precursors (1,4). Adipocytes are most abundant in adult human bone marrow, but there have been conflicting reports on their role in hematopoiesis (4). Adipocyte differentiation from murine preadipocytes has been shown to maintain hematopoiesis *in vitro* (5). In human long-term bone marrow culture, stromal adipocytes have been reported to be essential for long-term hematopoiesis, indicating that bone marrow adipocytes exert a positive effect on hematopoiesis (6). On the other hand, it has been shown that the ability of a preadipose cell line to support growth of hematopoietic stem cells decreases when the preadipocytes differentiate into adipocytes (7), and continuous human marrow hematopoiesis *in vitro* is not stimulated by adipocytes (8). Thus, the precise role of adipocytes in hematopoiesis remains to be determined.

Bone marrow stromal cells contain undifferentiated cells of mesenchymal origin which have the potential to differentiate into adipocytes, osteoblasts, chondrocytes, and stromal fibroblasts, which are referred to as mesenchymal stem cells (MSCs) (9). MSCs in bone marrow are few in number, but can be expanded *in vitro* to form a monolayer in culture of homogeneous populations. These cells have been shown to be able to support hematopoietic differentiation of CD34⁺ hematopoietic progenitor cells (10). It has also been reported that MSCs support the hematopoietic cells, including long-term culture initiating cells (LTC-IC) after differentiation toward stromal and osteogenic lineages (11).

In the present study, we compared the hematopoietic-supporting capacity of adipocytes and their progenitor, MSCs. We co-cultured cord blood (CB) CD34⁺CD38⁻ cells (an immature population of CD34⁺ cells) (12) on MSCs or adipocytes, and found that the hematopoiesis-supporting

Correspondence to: Dr Yoshihiro Fujimori, Laboratory of Cell Transplantation, Institute for Advanced Medical Sciences, Hyogo College of Medicine, 1-1 Mukogawa-cho, Nishinomiya, Hyogo 663-8501, Japan
E-mail: fuji-y@hyo-med.ac.jp

Key words: hematopoietic progenitor cells, CD34, mesenchymal stem cells, adipocytes, severe combined immunodeficiency repopulating cells

capacity of MSCs decreased with adipocyte differentiation. However, CB CD34⁺CD38⁻ cells co-cultured with adipocytes preserved their ability to engraft in NOD/SCID mice, suggesting that adipocytes maintain the ability to support transplantable SCID-repopulating cells.

Materials and methods

Cytokines. Recombinant human (rh)-interleukin (IL)-3, rh-stem cell factor (SCF), rh-granulocyte colony-stimulating factor (G-CSF), rh-granulocyte/macrophage (GM)-CSF, rh-thrombopoietin (TPO) and rh-erythropoietin (Epo) were a generous gift from the Kirin Brewery Co. Ltd. (Tokyo, Japan). Flt3 ligand (FL) was purchased from R&D Systems (Minneapolis, MN).

Mice. Ten-week-old female NOD/shi/SCID mice were purchased from Clea Japan (Tokyo, Japan). The mice were kept on racks under specific pathogen-free conditions with a laminar air flow and were supplied with sterile food and drinking water.

Culture of mesenchymal stem cells. Human bone marrow was aspirated from the iliac crest of normal subjects after obtaining informed consent. Mononuclear cells (MCs) were separated by density gradient centrifugation using Ficoll-Paque (Amersham Pharmacia, Uppsala, Sweden). MCs were cultured in a Mesenchymal stem cell growth medium (MSCGM) (Cambrex Biosciences, Walkersville, MD) (13). The medium was changed every week, and adherent cells with a fibroblast-like appearance were maintained. For adipogenic differentiation, MSCs were cultured in an adipogenic medium containing insulin, dexamethasone and indomethacin (hMSC differentiation kit-Adipogenic) (Cambrex Biosciences). For osteogenic differentiation, MSCs were incubated in an osteogenic medium containing dexamethasone, ascorbate and β -glycerolphosphate (Osteogenic differentiation kit). For chondrogenic differentiation, MSCs were cultured in a chondrogenic medium containing tumor growth factor (TGF) β -3, dexamethasone and ascorbate (Chondrogenic differentiation kit). Adipogenic, osteogenic and chondrogenic differentiation were confirmed by staining with Oil red O, von Kossa and Safranin O, respectively (14,15).

Isolation of lineage-negative cord blood cells. Umbilical cord blood (CB) was obtained from normal full-term deliveries after obtaining the mother's consent. Mononuclear cells (MCs) were separated by density gradient centrifugation using Ficoll-Paque (Amersham). The MCs were subjected to deplete lineage positive cells using the Human progenitor cell enrichment kit (StemCell Technologies, Inc., Canada) which included antibodies to lineage-specific antigens (CD2, CD3, CD14, CD16, CD19, CD24, CD56, CD66b, and Glycophorin A). The lineage negative cord blood cells were frozen in α -medium supplemented with 10% dimethylsulfoxide, 12% hydroxyethyl starch (CP-1 cryoprotectant) (Kyokuto Pharmaceutical Co., Tokyo, Japan) and 8% human serum albumin at -80°C.

Flow cytometric analysis and cell sorting. Cells were stained with fluorescein isothiocyanate (FITC)- or phycoerythrin

(PE)-conjugated monoclonal antibodies at 4°C for 30 min and subjected to an immunofluorescence analysis using an EPICS XL flow cytometer (Coulter, Miami, FL). The monoclonal antibodies used were PE-conjugated antibodies to CD34 (Coulter), CD45 (Coulter) and CD106 (Pharmingen, San Diego, CA); and FITC-conjugated antibodies to CD38 (Coulter), CD49, CD54 (Immunotech, Marceille, France), CD73 (Pharmingen) and CD105 (Serotec Oxford, UK). Dead cells were gated out with a forward vs side scatter window and propidium iodide staining. Isotype-matched antibodies were used as control in all experiments. CD34⁺CD38⁻ cells were isolated by sorting lineage negative cord blood cells using FACSaria (Becton-Dickinson).

Co-culture of CB CD34⁺CD38⁻ cells and MSCs or adipocytes in the presence of cytokines. A monolayer of MSCs was established in wells of a 12-well plate (Coster), and adipocytes were induced as described above. CD34⁺CD38⁻ cells (1×10^3 cells/well) were cultured without a stromal layer or on a layer of MSCs or adipocytes in 1 ml of serum-free medium, StemPro-34 (Gibco BRL, Grand Island, NY), supplemented with StemPro-34 nutrient supplement (Gibco BRL), 2 mM glutamine and penicillin/streptomycin in the presence of 100 ng/ml SCF, 100 ng/ml TPO and 100 ng/ml FL. After 7 and 14 days of culture, the cells were harvested by vigorous pipetting and subjected to CFC assay and flow cytometric analysis.

RT-PCR analysis of cytokines. Total RNA was extracted from the monolayer stromal cells using an Isogen RNA extraction kit (Wako Pure Chemical Co., Osaka, Japan) and was dissolved in diethylpyrocarbonate (DEPC; Sigma)-treated water. Reverse-transcriptase polymerase chain reaction (RT-PCR) was performed using primers described previously (13) and the RNA PCR kit AMV ver. 2.1 (Takara Shuzo, Shiga, Japan). The amplified products were separated by 2% agarose gel electrophoresis, stained with ethidium bromide, and photographed.

Colony-forming cell assay. Colony-forming cell (CFC) assays were performed in 35-mm petri dishes (Becton-Dickinson) by incubating the cells in semisolid α -medium containing 0.8% methylcellulose (Shinetsu Chemicals Co., Tokyo, Japan), 30% FCS (Gibco), 1% BSA, 10^{-4} M 2-mercaptoethanol (2-ME) (Wako Pure Chemicals), 2 mM L-glutamine (Sigma), 10 ng/ml IL-3, 20 ng/ml SCF, 10 ng/ml G-CSF, 10 ng/ml GM-CSF and 2 U/ml Epo (Kirin Brewery) for 14 days at 37°C in a humidified atmosphere flushed with 5% CO₂ in air (13). The colony-forming units (CFU)-GM, CFU-Mix and burst-forming units-erythroid (BFU-E) were identified by the ability to form granulocyte/macrophage (GM) colonies, mixed erythroid and myeloid (Mix) colonies and erythroid burst colonies, respectively, as described previously (16-18). High proliferative potential-colony-forming cells (HPP-CFC) were identified by their ability to form densely packed colonies that reached a size of >1 mm in diameter (19).

Transplantation into NOD/SCID mice (SCID-repopulating cell assay). CB CD34⁺CD38⁻ cells (1×10^3 cells/well in a 12-

Figure 1. MSCs and differentiation into adipocytes. (A) Phase microscopy of human mesenchymal stem cells (x40). (B) Flow cytometric analysis of cell surface markers on MSCs. Adipocytic differentiation of MSC was confirmed by the presence of neural lipid vacuoles (C) (x40) and lipid positive for Oil red O staining (D) (x40).

well plate) were cultured on a layer of MSCs or adipocytes or without a stromal layer in the presence of SCF/TPO/FL for 2 weeks. The cells harvested from one well were injected into the tail veins of 10-week-old NOD/shi/SCID mice sublethally irradiated (3Gy X-ray). The mice were intra-peritoneally injected with anti-asialo GM1 antibody (Wako Pure Chemical) to reduce the natural killer cell activity (20). The mice were sacrificed 8 weeks after the transplantation, DNA was extracted from BM cells using a SepaGene extraction kit (Sanko Pure Chemical Co., Tokyo, Japan) and was analyzed for the human-specific DNA 17 α -satellite gene by quantitative real-time PCR (21). The primers used were sense 5'-ACGGGATAACTGCACCTAAC-3', and antisense 5'-CCATAGGAGGGTTCAACTCT-3'.

Statistical analysis. The data are presented as the mean $\pm$ standard deviation (SD). The Student's t-test was used. P values of <0.05 were accepted as significant.

Results

Differentiation of MSCs into adipocyte lineage. Cultivation of bone marrow mononuclear cells from normal subjects in a mesenchymal stem cell growth medium (MSCGM) resulted in gradual proliferation of adherent MSCs. These cells were fibroblast-like in morphology (Fig. 1A), positive for CD73, CD105 (Fig. 1B), CD49, CD54, CD90 and CD106 (data not shown), and negative for CD34 and CD45 (Fig. 1B). Differentiation of MSCs into adipocytes was determined by the presence of neural lipid vacuoles (Fig. 1C) and lipid detectable by Oil red O stain (Fig. 1D). MSCs were also capable of undergoing osteogenic and chondrogenic differentiation under specific conditions (data not shown).

Figure 2. RT-PCR analysis of cytokine mRNAs. Total RNA was extracted from MSCs and adipocytes and analyzed for various cytokine mRNAs by RT-PCR.

The expression of cytokines and growth-related molecules in MSCs and adipocytes. We performed RT-PCR analysis of the mRNA of cytokines and growth-related molecules in MSCs and adipocytes. MSCs expressed SCF, TGF- β 1 and Jagged-1 but not TPO, IL-3 and GM-CSF (Fig. 2). The expression of SCF, TGF- β 1 and Jagged-1 was maintained after adipocyte differentiation. The message of adiponectin, an adipocyte-specific secretory protein (22), was not detected in MSCs but was expressed in adipocytes (Fig. 2).

Figure 3. Effect of adipocytes on the proliferation of CB34⁺38⁻ cells. CB CD34⁺CD38⁻ cells (1x10³) were cultured without stromal cells, or on MSCs or adipocytes in the presence of SCF, TPO and FL. On day 10 and 14, cells were collected and analyzed for the number of total viable cells (A) CD34⁺ cells (B) and CD34⁺38⁻ cells (C). Data are the mean $\pm$ SD of fold increases over the values on day 0 of four separate experiments. *P<0.05 in comparison to the MSC culture.

Level of engraftment (% human cells)

Primary CD34⁺ CD38⁻ cells

Cultured without stromal cells

Cultured with MSCs

Cultured with adipocytes

Group	Level of engraftment (% human cells)
Primary CD34 ⁺ CD38 ⁻ cells	~3.0
	~1.0
	~0.8
	~0.08
	~0.04
Cultured without stromal cells	~0.001
	~0.001
	~0.001
	~0.001
	~0.001
Cultured with MSCs	~0.15
	~0.08
	~0.04
	~0.02
	~0.015
Cultured with adipocytes	~0.25
	~0.1
	~0.02
	~0.01
	~0.008

performing SCID-repopulating cell (SRC) assays. All the progeny of CD34⁺CD38⁻ cells (1×10^3) cultured on MSCs or adipocytes in the presence of SCF/TPO/FL for 14 days were transplanted into NOD/SCID mice through the tail vein, and 8 weeks later, the mice were analyzed for human hematopoietic cells in the bone marrow by identifying the human-specific 17 α -satellite DNA using real-time PCR. All five mice transplanted with CB CD34⁺CD38⁻ cells cultured on MSCs were positive for human DNA, indicating that MSCs maintain SRCs (Fig. 5). All five mice transplanted with CB CD34⁺CD38⁻ cells cultured on adipocytes were also positive

Effect of MSCs or adipocytes on SCID-repopulating cells. We next investigated the ability of MSCs or adipocytes to maintain transplantable primitive progenitor cells by

for human DNA, thus indicating that adipocytes are also able to support SCID-repopulating cells. Control mice, transplanted with CB CD34⁺CD38⁻ cells cultured without stromal cells, were negative for the human DNA, indicating the loss of the SCID-repopulating capacity during liquid culture >9 days (23). These results show that adipocytes provide a milieu that maintains transplantable primitive hematopoietic progenitors.

Discussion

Adipocytes are the most abundant stromal cell type in adult human bone marrow, particularly in the long bone (4). In aplastic anemia patients, the bone marrow contains a reduced number of hematopoietic cells with abundant adipocytes. The increase in marrow fat in aplastic anemia is known to be caused by an increase in the size and number of individual fat cells. However, the precise role of adipocytes in the bone marrow is not fully understood. In the present study, we analyzed the hematopoietic-supporting capacity of adipocytes.

MSCs have the potential to differentiate into osteoblasts, adipocytes and stromal fibroblasts which constitute bone marrow stromal cells. MSCs have been shown to support hematopoiesis (10,11), but whether human marrow adipocytes possess a hematopoietic-supporting capacity remains controversial (6,8). It is also not clear whether marrow adipocytes are able to support SCID-repopulating cells, which are immature transplantable hematopoietic stem cells. Our results showed that bone marrow adipocytes were inferior to MSCs in supporting hematopoietic progenitors (Figs. 3 and 4). However, adipocytes were found to be similar to MSCs in maintaining SCID-repopulating cells (Fig. 5). Considering that only a small population of the CD34⁺CD38⁻ cells were SCID-repopulating cells, it is possible that adipocytes provided a niche to attach SCID-repopulating cells, resulting in their selective preservation.

The mechanism by which adipocytes regulate hematopoiesis remains unknown. When MSCs differentiated into adipocytes, adiponectin, an adipocyte-specific molecule, was expressed (Fig. 2). Adiponectin suppressed mature hematopoietic progenitors, such as CFU-GM, CFU-M and CFU-G, but not relatively immature progenitors, such as BFU-E and CFU-Mix (22). We observed that adipocytes, like MSCs, expressed mRNA of SCF, TGF- β 1 and Jagged-1 (Fig. 2). Notch-1 was expressed on CD34⁺ hematopoietic cells (24) and its ligand, Jagged-1, was expressed on stromal cells (25). Jagged-1 promoted hematopoiesis (25) and was able to expand SCID-repopulating cells (26). As adipocytes express such growth-related molecules, they may be able to provide an environment favorable for the maintenance of SCID-repopulating cells.

Our finding that adipocytes do not efficiently support mature hematopoietic progenitors may be consistent with the fact that mature blood cells, such as erythrocytes, WBC and platelets decrease in number in hypoplastic adipocyte-rich marrow in aplastic anemia. However, the hemopoietic micro-environment in adipocyte-rich marrow may still be able to preserve immature hematopoietic progenitor cells. Thus, adipocytes in the bone marrow may not be simple replacements of the marrow cavity but may positively influence immature hematopoietic progenitor cells.

Acknowledgements

We are grateful to Ms. Ikuyo Kasumoto and Ms. Hiromi Takeda for their excellent technical assistance. This study was supported by a research grant for the 'High-Tech Research Center' Project for Private Universities from the Ministry of Education, Culture, Sports, Science and Technology of Japan, and a grant from the Research on Human Genome, Tissue Engineering Food Biotechnology of the Ministry of Health, Labor and Welfare of Japan.

References

1. Dorshkind K: Regulation of hematopoiesis by bone marrow stromal cells and their products. *Annu Rev Immunol* 8: 111-137, 1990.
2. Baines P, Mayani H, Bains M, Fisher J, Hoy T and Jacobs A: Enrichment of CD34 (My10)-positive myeloid and erythroid progenitors from human marrow and their growth in cultures supplemented with recombinant human granulocyte-macrophage colony-stimulating factor. *Exp Hematol* 16: 785-789, 1988.
3. Larochelle A, Vormoor J, Hanenberg H, Wang JC, Bhatia M, Lapidot T, Moritz T, Murdoch B, Xiao XL, Kato I, Williams DA and Dick JE: Identification of primitive human hematopoietic cells capable of repopulating NOD/SCID mouse bone marrow: implications for gene therapy. *Nat Med* 2: 1329-1337, 1996.
4. Gimble JM, Robinson CE, Wu X and Kelly KA: The function of adipocytes in the bone marrow stroma: an update. *Bone* 19: 421-428, 1996.
5. Greenberger JS: Sensitivity of corticosteroid-dependent insulin-resistant lipogenesis in marrow preadipocytes of obese-diabetic (db/db) mice. *Nature* 275: 752-754, 1978.
6. Gartner S and Kaplan HS: Long-term culture of human bone marrow cells. *Proc Natl Acad Sci USA* 77: 4756-4759, 1980.
7. Kodama H, Sudo H, Koyama H, Kasai S and Yamamoto S: *In vitro* hemopoiesis within a microenvironment created by MC3T3-G2/PA6 preadipocytes. *J Cell Physiol* 118: 233-240, 1984.
8. Touw I and Lowenberg B: No stimulative effect of adipocytes on hematopoiesis in long-term human bone marrow cultures. *Blood* 61: 770-774, 1983.
9. Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, Mosca JD, Moorman MA, Simonetti DW, Craig S and Marshak DR: Multilineage potential of adult human mesenchymal stem cells. *Science* 284: 143-147, 1999.
10. Majumdar MK, Thiede MA, Mosca JD, Moorman M and Gerson SL: Phenotypic and functional comparison of cultures of marrow-derived mesenchymal stem cells (MSCs) and stromal cells. *J Cell Physiol* 176: 57-66, 1998.
11. Majumdar MK, Thiede MA, Haynesworth SE, Bruder SP and Gerson SL: Human marrow-derived mesenchymal stem cells (MSCs) express hematopoietic cytokines and support long-term hematopoiesis when differentiated toward stromal and osteogenic lineages. *J Hematother Stem Cell Res* 9: 841-848, 2000.
12. Petzer AL, Zandstra PW, Piret JM and Eaves CJ: Differential cytokine effects on primitive (CD34⁺CD38⁻) human hematopoietic cells: novel responses to Flt3-ligand and thrombopoietin. *J Exp Med* 183: 2551-2558, 1996.
13. Nishioka K, Fujimori Y, Hashimoto-Tamaoki T, Kai S, Qiu H, Kobayashi N, Tanaka N, Westerman KA, Leboulch P and Hara H: Immortalization of bone marrow-derived human mesenchymal stem cells by removable simian virus 40T antigen gene: analysis of the ability to support expansion of cord blood hematopoietic progenitor cells. *Int J Oncol* 23: 925-932, 2003.
14. Jaiswal N, Haynesworth SE, Caplan AI and Bruder SP: Osteogenic differentiation of purified, culture-expanded human mesenchymal stem cells *in vitro*. *J Cell Biochem* 64: 295-312, 1997.
15. Mackay AM, Beck SC, Murphy JM, Barry FP, Chichester CO and Pittenger MF: Chondrogenic differentiation of cultured human mesenchymal stem cells from marrow. *Tissue Eng* 4: 415-428, 1998.
16. Hara H, Kai S, Fushimi M, Taniwaki S, Okamoto T, Ohe Y, Fujita S, Noguchi K, Senba M, Hamano T, Kanamaru A and Nagai K: Pluripotent hemopoietic precursors *in vitro* (CFUMIX) in aplastic anemia. *Exp Hematol* 8: 1165-1171, 1980.

17. Fujimori Y, Hara H and Nagai K: Effect of lymphokine-activated killer cell fraction on the development of human hematopoietic progenitor cells. *Cancer Res* 48: 534-538, 1988.
18. Fujimori Y, Ogawa M, Clark SC and Dover GJ: Serum-free culture of enriched hematopoietic progenitors reflects physiologic levels of fetal hemoglobin biosynthesis. *Blood* 75: 1718-1722, 1990.
19. McNiece IK, Stewart FM, Deacon DM, Temeles DS, Zsebo KM, Clark SC and Quesenberry PJ: Detection of a human CFC with a high proliferative potential. *Blood* 74: 609-612, 1989.
20. Yoshino H, Ueda T, Kawahata M, Kobayashi K, Ebihara Y, Manabe A, Tanaka R, Ito M, Asano S, Nakahata T and Tsuji K: Natural killer cell depletion by anti-asialo GM1 antiserum treatment enhances human hematopoietic stem cell engraftment in NOD/Shi-scid mice. *Bone Marrow Transplant* 26: 1211-1216, 2000.
21. Becker M, Nitsche A, Neumann C, Aumann J, Junghahn I and Fichtner I: Sensitive PCR method for the detection and real-time quantification of human cells in xenotransplantation systems. *Br J Cancer* 87: 1328-1335, 2002.
22. Yokota T, Oritani K, Takahashi I, Ishikawa J, Matsuyama A, Ouchi N, Kihara S, Funahashi T, Tenner AJ, Tomiyama T and Matsuzawa Y: Adiponectin, a new member of the family of soluble defense collagens, negatively regulates the growth of myelomonocytic progenitors and the functions of macrophages. *Blood* 96: 1723-1732, 2000.
23. Bhatia M, Bonnet D, Kapp U, Wang JC, Murdoch B and Dick JE: Quantitative analysis reveals expansion of human hematopoietic repopulating cells after short-term *ex vivo* culture. *J Exp Med* 186: 619-624, 1997.
24. Milner LA, Kopan R, Martin DI and Bernstein ID: A human homologue of the *Drosophila* developmental gene, Notch, is expressed in CD34+ hematopoietic precursors. *Blood* 83: 2057-2062, 1994.
25. Jones P, May G, Healy L, Brown J, Hoyne G, Delassus S and Enver T: Stromal expression of Jagged 1 promotes colony formation by fetal hematopoietic progenitor cells. *Blood* 92: 1505-1511, 1998.
26. Karanu FN, Murdoch B, Gallacher L, Wu DM, Koremoto M, Sakano S and Bhatia M: The notch ligand jagged-1 represents a novel growth factor of human hematopoietic stem cells. *J Exp Med* 192: 1365-1372, 2000.